

# Grandmothers and the Evolution of Human Longevity

KRISTEN HAWKES\*

*Department of Anthropology, University of Utah, Salt Lake City, Utah 84112*

**ABSTRACT** Great apes, our closest living relatives, live longer and mature later than most other mammals and modern humans are even later-maturing and potentially longer-lived. Evolutionary life-history theory seeks to explain cross-species differences in these variables and the covariation between them. That provides the foundation for a hypothesis that a novel role for grandmothers underlies the shift from an ape-like ancestral pattern to one more like our own in the first widely successful members of genus *Homo*. This hypothesis links four distinctive features of human life histories: 1) our potential longevity, 2) our late maturity, 3) our midlife menopause, and 4) our early weaning with next offspring produced before the previous infant can feed itself. I discuss the problem, then, using modern humans and chimpanzees to represent, respectively, genus *Homo* and australopithecines, I focus on two corollaries of this grandmother hypothesis: 1) that ancestral age-specific fertility declines persisted in our genus, while 2) senescence in other aspects of physiological performance slowed down. The data are scanty but they illustrate similarities in age-specific fertility decline and differences in somatic durability that are consistent with the hypothesis that increased longevity in our genus is a legacy of the “reproductive” role of ancestral grandmothers. *Am. J. Hum. Biol.* 15:380–400, 2003. © 2003 Wiley-Liss, Inc.

Based on detailed examination of human reproductive aging, Gosden (1996:281) concluded that the “reproductive system ages faster than the body as a whole and by age 45 can be said to be in the state that a woman’s other organs have reached by eighty.” Here I review the foundation for linking that characteristic of humans to other life-history features that distinguish us from our nearest living relatives, the chimpanzees. After briefly outlining the problem, I consider the hunting hypothesis, an influential explanation for some of those features based on benefits from large-game hunting among our ancestors. Summarizing archaeological and paleontological evidence against the hunting hypothesis, I present a simplified outline of hominin evolution that highlights the life-history shift in genus *Homo*. Then I review ideas from evolutionary life-history theory and cross-species variations they explain. Theory and data linking adult mortality risk to both maturation and aging rates provide the foundation for a grandmother hypothesis to account for the life-history shift in our genus. I reprise that hypothesis and then report comparisons of age-specific changes in fertility, mortality, and aging between humans and chimpanzees. These data, while consistent with the hypothesis, are very limited, highlighting how much more we need to know about fertility and aging in chimpanzees.

## PROBLEMS OF HUMAN LIFE HISTORY

In three of the classic articles that established evolutionary life-history theory (Cole, 1954; Williams, 1957; Hamilton, 1966), long postmenopausal survival was recognized as a striking characteristic of modern humans. Williams (1957:407), noting that selection could maintain “little or no post-reproductive period in the normal life-cycle of any species” proposed the most influential hypothesis to explain the apparent exception in humans: “At some time during human evolution it may have become advantageous for a woman of forty-five or fifty to stop dividing her declining faculties between the care of extant offspring and the production of new ones.” This scenario implies that modern aging rates were established in ancestral human populations before subsequent adjustment in our age at menopause. Analytical attention thus focused on the optimal timing of fertility termination in humans (e.g., Hawkes et al., 1989; Hill and Hurtado, 1991, 1996, 1999; Rogers, 1993; Pectei, 2001; Shanley and Kirkwood, 2001),

\*Correspondence to: Kristen Hawkes, Anthropology, 270 E 1400 S Stewart 102, University of Utah, Salt Lake City UT 84112-0600. E-mail: hawkes@anthro.utah.edu

Received 16 August 2002; Accepted 9 December 2002

Published online in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/ajhb.10156

and comparisons among the lengths of post-fertile lifespans across species (e.g., Finch, 1990; Pavelka and Fedigan, 1991; vom Saal et al., 1994; Caro et al., 1995; Packer et al., 1998).

But since Williams proposed this stopping early hypothesis, theoretical and empirical reasons have been accumulating to pose questions about human postmenopausal longevity in the opposite way. Our age at menopause may be the conserved ancestral trait, while our long potential lifespans are derived (Kaplan, 1997; Judge and Carey, 2000). Cross-species variation in longevity and links between that variable and other life-history features then become matters of central interest. Maximum lifespans vary more than 50-fold among mammalian species. That variation is positively correlated with differences in maturation rates, a relationship that persists even when the body size effects are removed (Harvey and Zammuto, 1985; Sutherland et al., 1986; Harvey et al., 1989; Read and Harvey, 1989; Promislow et al., 1990). Theory to explain the cross-species regularities (Charnov, 1991, 1993, 1997; Charnov and Berrigan, 1991) combined with a hypothesis about a novel role for ancestral grandmothers (Hawkes et al., 1998) and evolutionary theories of aging (Williams, 1957; Hamilton, 1966; Kirkwood, 1977; Ricklefs, 1998) can account for a shift toward greater longevity and later maturity, while ancestral age-specific fertility termination was conserved in our lineage. This hypothesis highlights ancestral socioecological circumstances that allowed more vigorous perimenopausal females to raise the fertility of their daughters, consequently strengthening selection against senescence (the greater vulnerability to mortality with increasing age) and increasing potential longevity in our genus.

Using a demographic approach to life histories, the hypothesis relies on models in which adult mortality risk determines selection on both age at maturity and rates of senescence. Theory and evidence highlight the increase in intrinsic (senescent) components of adult mortality risk when life histories are slow (Williams, 1957; Ricklefs, 1998). This is justification for the proposition that reduced aging rates can further lessen adult mortality risk in already slow life histories. The demographic approach contrasts with others that use variation in fetal, infant, and juvenile patterns of growth and development to explain matur-

ation rates (e.g., Gould, 1977; Leigh, 2001; Lieberman, 2002). Starting with adult mortality reveals cross-species regularities that are extremely general and this approach to human life histories links four distinctive features to each other: our long potential lifespans, late maturity, short interbirth intervals, and midlife menopause.

The most influential and most fully elaborated alternative argument to explain the evolution of late maturity and high fertility in humans, the hunting hypothesis, proposes that these features are consequences of expanding human brains. I outline it and then sketch the current paleoanthropological record of the timing and character of transitions in human evolution. The summary highlights both lack of fit with the hunting hypothesis and the prominence of a life-history shift in the evolution of our genus. Then I assemble the combination of theory and evidence that justifies a grandmother hypothesis to explain the evolution of genus *Homo* and proceed to comparisons between humans and chimpanzees that help to reveal our design for fertility and longevity.

#### THE HUNTING HYPOTHESIS

The hunting hypothesis was the most influential argument about human origins throughout the second half of the last century (Dart, 1953; Washburn, 1960; Washburn and DeVore, 1961; Washburn and Lancaster, 1968; Isaac, 1978; Hill, 1982; Lancaster and Lancaster, 1983, 1987; Tooby and DeVore, 1987; Kaplan et al., 2000). There are good reasons for the persistence of this idea. Cartmill (1993:191), reviewing its history, noted that "the hunting hypothesis was the first truly Darwinian explanation of human origins to be proposed." Building on Dart's proposals (e.g., Dart, 1949, 1953), it was Washburn who articulated the hunting hypothesis most effectively within anthropology (e.g., Washburn and Avis, 1958; Washburn and DeVore, 1961). In this hypothesis, features of life history and social organization that distinguish humans from other primates evolved because increased reliance on hunting large animal prey gave net advantages to larger brains. Its main elements are these:

- Drying environments in the late Tertiary constricted African forests, making

capacities to use alternative foods more advantageous among ancestral apes.

- Bipedalism was then favored because it freed hands for tool use, which increased success at hunting big animals, and this put a premium on larger brains.
- But the mechanics of bipedal locomotion limited pelvic width, so brain expansion created an “obstetrical dilemma” requiring most brain growth to be postnatal. Consequently, children with developing brains were immature longer and were more dependent, for a longer time, on maternal care.
- The care requirements interfered with maternal hunting, so mothers relied on provisioning from hunting mates. This help from fathers allowed mothers to produce more surviving offspring.
- Thus, parents formed lasting bonds and nuclear families became the fundamental units of cooperation in which a sexual division of labor served familial goals of production and reproduction.

This hypothesis was especially influential because it tied together the earliest archaeology, the fossil evidence of human ancestry, and social organization differences between humans and other primates. Washburn championed investigation of all those lines of evidence.

### THE EARLY ARCHAEOLOGY

Paleolithic archaeologists used the hunting hypothesis to interpret the rich east African PlioPleistocene sites that are roughly the same age as the first fossils assigned to our genus. Isaac (1978) saw these as living areas where ancient hunters brought their prey to provision families. This view was subsequently challenged on grounds that the bone assemblages were more likely the residue of hominin scavenging from remains abandoned after most meat was consumed by the primary carnivore predators (Binford, 1981; Blumenshine, 1987). Countering this, other analysts found cut marks on bones indicating that the hominins also had access to carcasses that were still well fleshed (Bunn and Ezzo, 1993). Some archaeologists interpret the damage patterns as plausible indications of hunting (Dominguez-Roderigo, 2002). O’Connell et al. (2002a,b) note the locations of these early

sites to be likely settings for carnivores to ambush prey. They review the mixed patterns of tool and carnivore tooth marks on the bones and surmise variable hominin success at competitive scavenging. Using modern analogs to estimate the likely amount of meat hominins might procure in this way, they show amounts too variable to be a major source of nutrition (O’Connell et al., 2002a,b). Since Washburn’s elaboration of the hunting hypothesis stimulated detailed examination of the early archaeology (Issac, 1984), consensus that it indicates big-game hunting by the first members of our genus has collapsed.

### HUMAN PALEONTOLOGY

The fossil evidence of the hominins themselves has expanded substantially in the last three decades. Evidence that bipedalism was not linked to brain expansion came first (Johanson and White, 1979). Now three main radiations of fossil hominins can be (provisionally) distinguished in the record (Klein, 1999, 2000). First was the initial divergence of our clade from that of chimpanzees, marked by a shift to bipedality, perhaps around 6 mya (e.g., Haile-Selassie, 2001; Senut et al., 2001). The brain and body sizes of those australopithecines (used in the broad sense to include *Australopithecus*, *Ardipithecus*, *Kenyanthropus*, *Paranthropus*, etc.) were similar to modern chimpanzees (McHenry, 1994). Their maturation patterns were more similar to modern chimpanzees than to modern humans (Bromage and Dean, 1985; Smith, 1986, 1987, 1989, 1991, 1994; Smith and Tompkins, 1995; Dean et al., 2001). On those grounds, the several taxa in this australopithecine radiation have been characterized as “bipedal apes” (Klein, 1999). Bipedalism antedated both brain expansion and the archaeological record by millions of years.

The second radiation began around two million years ago with genus *Homo*. The first widely successful members of our genus, labeled *H. ergaster* in Africa (Wood, 1992) and *H. erectus* in Asia (Rightmire, 1990) showed a marked increase in brain size. Body sizes also increased substantially, making the *relative* brain size change less dramatic (McHenry, 1992, 1994; Aiello and Wheeler, 1995; Aiello and Wells, 2002; Wood

and Collard, 1999). Collard and Wood (1999: 324) go so far as to say that:

Although there are twofold differences in the mean absolute brain size of early hominids, these differences are almost certainly not significant when body mass is taken into account. A notable effect of body-mass correction is that the absolutely larger brain of *H. ergaster* is "cancelled out" by its substantial estimated body mass.

*Homo ergaster/erectus* reached its larger size by maturing later than the australopithecines (Smith BB, 1991, 1993, 1994; Dean et al., 2001; Aiello and Wells, 2002). With body size and shape similar to modern humans, maturation patterns may also have been within the modern range (Smith, 1994; Clegg and Aiello, 1999; Tardieu, 1998; Anton, 2001).

*Homo ergaster/erectus* had smaller teeth and jaws and a narrower thorax than australopithecines, indicating reliance on foods that require less mastication and digestion (Aiello and Wheeler, 1995). Different foraging strategies are also indicated by expansion into a wide range of new habitats. This taxon is the first hominin found outside Africa, as far east as Indonesia and as far north as 45° (Gabunia et al., 2000; Larick et al., 2001; Swisher et al., 1994; Zhu et al., 2001). Both the smaller teeth and guts and the range expansion have been attributed to increased meat eating (e.g., Shipman, 1986; Shipman and Walker, 1989; Milton, 1999; Stanford and Bunn, 2001), an inference often linked to the assumption that the PlioPleistocene sites represent big-game hunting. As noted above, archaeological challenges to that claim are numerous. Big-game hunting is not reliably indicated in the archeology until the last half million years (Stiner, 2002). Isotope analyses of a sample of specimens show no more meat eating in *H. ergaster* than among contemporaneous australopithecines (Lee-Thorp et al., 2000; Lee-Thorp, 2002). Alternatives (Wrangham et al., 1999), including the grandmother hypothesis (O'Connell et al., 1999, 2002a,b), may better explain the morphological changes.

It was with the third hominin radiation, that of archaic humans which began about 0.5 mya, that brains reached the large size of

moderns (e.g., McHenry, 1994; Ruff et al., 1997). So by this reading of the evidence, the second radiation, the appearance of genus *Homo*, is associated especially with a change in life history as indexed by body size and maturation rate and a change in resource use inferred from dentition, body form, and geographical range. Bipedalism precedes this life-history shift by millions of years. Big-game hunting and the very big brains that evolve with archaic humans (including both early near moderns and their contemporaries like the Neanderthals) do not appear until more than 1.5 million years later. The features linked in the hunting hypothesis do not evolve together and the large brains that are presumed to drive the evolution of other features emerge last.

Since Washburn elaborated the hunting hypothesis, not only the archeological and fossil evidence has changed. Theory and models unavailable then have become standard tools of analysis in behavioral evolution. Conflicts of interest among individuals have been recognized as especially important (Williams, 1966a; Trivers, 1972; Hrdy, 1981, 1999; Haig, 1993). Investigations into social organization in other primates reveal persistent male-female relationships that do not depend on economic cooperation (Smuts, 1985, 1992; van Schaik, 1996; van Schaik and Janson, 2000). Studies of many living populations now challenge previous assumptions that nuclear families evolve and persist as units of common reproductive interest (Gowaty, 1996; Mesnick, 1997; Palombit, 1998; Bliege Bird, 1999; Hawkes et al., 2001).

#### LIFE-HISTORY THEORY

At the same time that evolutionary anthropologists were especially stimulated by Washburn's extensions of the hunting hypothesis, MacArthur and Wilson (1967; Pianka, 1970) proposed r and K selection to account for the enormous interspecific variation in rates of maturation and offspring production among living things. They hypothesized that life-history tactics giving high maximal rates of population increase (r) would be favored when ecological disruptions result in population crashes and periodic opportunities for rapid population growth. These circumstances would favor early maturity at small size, with all effort expended in producing many small offspring, and so early death. On the other

hand, in “saturated” environments, with population densities near carrying capacity ( $K$ ), they hypothesized that selection would favor life histories that maximize competitive capacities: late maturity at large size and the production of a few large, well-developed offspring over long adult lives.

While the logic of  $r$  &  $K$  selection is simple and compelling, both empirical and theoretical work has shown that density-dependent mortality has different effects on developmental strategies depending on which age classes suffer the mortality (Gadgil and Bossert, 1970; Charnov and Shaffer, 1973; Stearns, 1977). The ecological associations originally postulated for  $r$  &  $K$  selection do not hold empirically. Both clusters of characters can evolve and persist in stable populations (see Stearns, 1992, for review). But the fast/slow variation that was of initial interest (MacArthur and Wilson, 1967; Pianka, 1970) and correlations between life-history variables and body size (Bonner, 1965; Blueweiss et al., 1978; Western, 1979; Harvey and Clutton-Brock, 1985) are real regularities of the empirical world.

#### *Adult lifespans and age at maturity*

Life-history features are not only correlated with body size, they also correlate with each other when the effects of body size are removed (Harvey and Read, 1988; Read and Harvey, 1989; Promislow et al., 1989; Harvey et al., 1989). During the last decade, Charnov, building on his own earlier work and that of many others (e.g., Koslowski and Wiegert, 1987; Harvey and Purvis, 1999, for review), focused on only a very few tradeoffs that can account for these patterns. Within taxonomic groups, larger-bodied species live slower lives, but some taxa are relatively slower, and others faster, at similar body sizes. The explanatory models take account of both the within-taxon correlations with body size and the between-taxon differences (Charnov, 1991, 1993, 1997, 2001; Charnov and Berrigan, 1991; Purvis and Harvey, 1995).

Charnov’s models capture both similarities in the relationships among life-history features in primate and nonprimate mammals and also some distinctive features of the primate order (Charnov, 1993; Charnov and Berrigan, 1993). The “mouse lemur to gorilla” curve describes allometric variation in the life-history variables with body size

that parallels the “mouse to elephant” curve. In both cases larger bodies are associated with longer adult lifetimes (the inverse of the adult mortality rate:  $M$ ), later ages at maturity ( $\alpha$ ), and lower annual fecundity ( $b$ ). These features are not only correlated within each group, Charnov (1991, 1993) shows that their products are “invariant” even as the values of the variables themselves change across the range of body sizes.

Primates are slower than nonprimate mammals, taking longer to mature at a given size and reproducing more slowly for size, but relationships among the life-history variables are the same for primate and nonprimate mammals. Charnov explains the relationships using a simple growth model in which productive capacity is an allometric function of size. [Charnov (2001) modifies this assumption to allow more realistic sigmoid growth, a modification that does not change the basic results discussed here.] Juveniles are assumed to use their productive capacity to grow themselves, then, at maturity, adults redirect that capacity into producing offspring. The benefit of delaying reproduction is the greater productive capacity resulting from growth to larger size. The cost of delaying maturity is the risk of dying before reproducing. If adult mortality risk is high, selection favors maturing earlier. If adult mortality risk declines, selection favors growing longer before maturing to reap the productive benefits of larger size. Whether life histories slow down or speed up depends on adult mortality risk.

#### *Adult lifespans and rates of aging*

Adult mortality risk not only provides a key to variation in age at maturity and rates of offspring production, it is central to evolutionary explanations of senescence, defined as age-specific declines in adaptive performance (Williams 1957). Williams (1957:404) used the decline of selection pressures with increasing age to explain why “*low adult death rates should be associated with low rates of senescence, and high adult death rates with high rates of senescence . . . [so] we should be able to predict rates of senescence on the basis of adult mortality rates*” [original italics]. Paradoxically, senescence is faster in species with fast life histories because individuals do not live long enough to grow old. When adult mortalities

are low, more individuals live long enough to display age-specific declines in performance; selection against senescence strengthens; aging slows. But age-specific mortalities are more strongly affected by senescence the lower the adult mortalities. An increasingly larger fraction of adult mortality is due to age-specific frailty. Ricklefs (1998) demonstrates how sharply this intrinsic component of adult mortality increases with lifespan in mammals and birds.

Fitting a parameter for age-specific changes in adult mortality rates to data on 18 species of birds and 27 species of mammals, Ricklefs (1998) found that, as predicted, species with longer adult lifespans age more slowly. He used the difference between the minimum mortality rate (the rate in young adults) and rates at older ages to estimate intrinsic (senescent) mortality, and measured the fraction of adult deaths due to senescence for different lifespans. In populations with average adult life spans just under 2 years, less than 8% of adult deaths are due to senescence, i.e., all but those few deaths would occur at the observed ages even if there were no senescence and vulnerability to mortality remained constant with age. On the other hand, in populations with average adult life spans just over 15 years (the range for modern great apes), 69% of the adult deaths were due to senescence. Selection can only slow senescence to the extent that the benefits of the physiological adjustments that retard aging outweigh the cost. Very high proportions of senescent deaths in long-lived organisms indicate that these costs are considerable (Ricklefs, 1998).

#### *Disposable somas*

Life-history theory assumes that the allocation of energy and time to somatic durability and maintenance is limited by fundamental tradeoffs. Kirkwood's "disposable soma" model (Kirkwood, 1977, 1981; Kirkwood and Holliday, 1979; Kirkwood and Rose, 1991) is a version of Williams' (1957) antagonistic pleiotropy theory of aging that highlights the tradeoff between somatic effort and current reproductive effort. Selection favors disposable somas rather than immortal ones because physiological adjustments that slow senescence leave less for current reproduction. Lower reproductive rates cannot be favored without other compensating effects on lifetime fitness.

The tradeoff is demonstrated in experiments on *Drosophila*, where artificial selection can increase lifespans over only a few generations and increased longevity is accompanied by decreases in reproduction (Rose, 1991). Heritable variation that could allow selection to shift longevity quickly is widely distributed. Both pedigree analysis and selection experiments indicate substantial within-population variation in longevity in a wide array of species, including humans (Herskind et al., 1996; Finch and Tanzi, 1997; Carey and Judge, 2001). Tradeoffs between longevity and fertility appear in human datasets as well (e.g., Westendorp and Kirkwood, 1998).

Many physiological processes contribute to somatic durability, processes that operate not only in adulthood, but throughout life (Service et al., 1985). Reviewing the history of ideas, Holliday (1995:102) noted that:

Initially, the disposable soma theory took into account accuracy in macromolecular synthesis, ... [Then] the metabolic cost of repair of macromolecules was an obvious inclusion (Kirkwood, 1981), and later on many other types of mechanism were discussed in terms of the maintenance of the adult organism. ... Today the disposable soma theory includes the considerable metabolic expense of all such maintenance mechanisms and the tradeoff between this expense and the investment of resources into growth to adulthood and reproduction.

By focusing attention on these mechanisms, the disposable soma model reverses the question about variation in rates of senescence from "Why do organisms get old?" to the converse: "Why do organisms live so long?" Living processes result in damage that can be reduced or repaired, but always at some cost. Sometimes increased somatic maintenance and repair results in longer-term fitness benefits that outweigh those costs, including the cost of reduced, or postponed, reproduction (Williams, 1966b; Charnov and Schaffer, 1973; Stearns, 1992). Selection can therefore favor increased longevity, with lower rates of annual fecundity, as long as the overall rate of increase in the longer-lived lineages is higher than in competing lineages. The enormous empirical

variation in lifespans (Finch, 1990, 1997) and the generally lower rates of annual fecundity in longer-lived mammals and birds (Read and Harvey, 1988; Saether, 1988; Stearns, 1992; Charnov, 1993) are consistent with these tradeoff arguments.

#### GRANDMOTHER HYPOTHESIS AND THE LIFE-HISTORY SHIFT IN GENUS *HOMO*

Consider the evidence of human phylogeny in light of this broad and systematic variation in mammalian life histories. If australopithecines matured at about the same age as chimpanzees, implications follow for other aspects of their life histories. Modern chimpanzees mature very late and have maximum lifespans that are very long compared to other terrestrial mammals (Eisenberg, 1981). As noted above, *H. erectus* was larger, maturing later than the australopithecines (Smith, 1991, 1994; Clegg and Aiello, 1999; Tardieu, 1998; Anton, 2001; Dean et al., 2001). From a demographic perspective, later maturity implies even lower adult mortality and longer adult life spans than that typical of chimpanzees. The link between age at maturity and adult mortality poses the following question for the evolution of genus *Homo*: How could adult mortalities, already quite low in australopithecines, have been reduced even further in our genus?

#### *Ethnographic clues*

Observations among Hadza hunter-gatherers in northern Tanzania (Woodburn, 1968; Blurton Jones et al., 1992) suggest an answer. Among these modern people, women past menopause are productive and energetic foragers (Hawkes et al., 1989), and so are children (Blurton Jones et al., 1989). But young children are not very effective at handling an important diet staple, the large root of a plant that is deeply buried and requires some strength to excavate (Hawkes et al., 1995). Women past menopause are especially active tuber diggers (Hawkes et al., 1989). In this population, mothers' foraging effort has a measurable effect on their children's nutritional welfare except when those mothers are nursing new infants. Lactating women spend less time foraging, and it is the work of postmenopausal grandmothers that differentially affects the nutritional welfare of weaned children (Hawkes et al., 1997). Grandmaternal

effects on the welfare of weanlings have been found in some other contemporary and historical populations (e.g., Sear et al., 2000; Volland and Biese, 2002; Jamison et al., 2002), but not all (Hill and Hurtado, 1991, 1996, 1999). The ecological circumstances faced by Hadza foragers highlight an especially important fitness opportunity for older females that would have arisen with the past climate changes that Washburn incorporated into the hunting hypothesis.

Drying environments in the Pliocene constricted the availability of foods that young juveniles can handle. Increasing aridity and seasonality favor plants that cope well with dry seasons, for example, by holding nutrients in hard-cased seeds, nuts, and underground storage organs. Such resources can give high return rates to human foragers, but only to those with the strength and skill to extract and process them. Young juveniles cannot do it. To rely on these resources and succeed in these environments, mothers have to provision offspring who are still too young to extract and process the foods themselves. The mother-offspring provisioning allows the occupation of otherwise inhospitable environments. It also creates a novel fitness opportunity for older females whose own fertility is declining. If the older females help feed their just-weaned grandchildren, the mothers of those weanlings can have shorter interbirth intervals without reductions in offspring survivorship. The more vigorous elders who have no nursing infants of their own will thus raise their daughters' reproductive success (Hawkes et al., 1998; O'Connell et al., 1999).

#### *Grandmothers and Charnov's invariants*

This grandmother hypothesis combined with Charnov's model implies particular relationships among life-history variables. If Charnov's tradeoff arguments apply, and if females past childbearing are contributing to their daughters' reproductive success, then human age at maturity ( $\alpha$ ) should be adjusted to the whole adult lifespan ( $1/M$ , the inverse of the adult mortality rate), preserving the relationship between  $\alpha$  and  $M$  that is generally characteristic of primates. On the other hand, with grandmothers' help, human interbirth intervals during the childbearing years should be shorter than expected for a grandmother-less ape with our age at maturity.

In his 1993 book, Charnov displayed the correlation between age at maturity and average adult lifespan for 15 primate sub-families. His analysis showed the relationship to be no different from the relationship between these variables in nonprimate mammals. Charnov's (1993) graph showing this pattern in primates included a data point for hominins, which comes from the only living member of the taxon: modern humans. While both lifespan and age at maturity are much higher in modern humans than in other primates, the human point falls just as predicted by the general primate  $\alpha M$  invariant. As expected from this model, the  $\alpha M$  numbers for three other living great apes, orangutans, gorillas, and chimpanzees, are similar to each other and also to an  $\alpha M$  number calculated for a sample of modern human foragers (Hawkes et al., 1998). Alvarez (2000) confirms that the relationship between  $\alpha$  and  $M$  in humans is well within the confidence interval of the general primate pattern based on data from 16 primate species. The fact that living primates *including modern humans* illustrate this  $\alpha M$  invariance provides justification for the working assumption that the same relationships among life-history variables, and so the same fundamental tradeoffs, applied to ancestral hominins. If so, the ages at maturity of the fossil taxa imply characteristic longevities.

In addition to the  $\alpha M$  invariant, Charnov's model highlights an invariant relationship between age at maturity ( $\alpha$ ) and average annual fecundity ( $b$ ). For mammals generally and primates in particular, annual fecundity goes down as age at maturity is delayed. Later-maturing mothers are larger and nurse their babies to larger size before weaning. Consequently, they produce offspring at a slower rate. The grandmother hypothesis proposes that specific socioecological circumstances allow more vigorous elders to help their daughters provision weanlings, allowing those daughters to wean offspring earlier than without such help (Bogin, 1999, 2001). Consequently, selection favors increased vigor when fertility is dropping because the fecundity of the child-bearers in a grandmothere lineage is higher than expected for a grandmother-less one. Comparisons among orangutans, gorillas, chimpanzees, and humans show this to be the case (Hawkes et al., 1998). Alvarez (2000) confirmed that, as predicted by this

hypothesis, the human rate of offspring production is above the confidence intervals determined by the regular relationship between age at maturity ( $\alpha$ ) and annual fecundity ( $b$ ) across 16 primate species.

This apparent break with Charnov's "assembly rules" for mammalian life histories is, at the same time, consistent in a deeper sense with the general tradeoffs his model highlights. Humans have a higher probability of birth per year ( $b$ ) than expected in the childbearing years, with  $b$  falling to zero a few years before menopause. The argument here proposes that grandmothere results in two interdependent components of adulthood, with higher than expected fecundity in one *because* there is no direct fecundity in the other. A grandmother-less primate, modeled according to these tradeoffs to have our age at maturity, would have constant fecundity throughout adult life instead. When human fecundity is calculated as an average over all adulthood, not just the childbearing years, our  $\alpha b$  does approach the primate invariant.

#### *Grandmothers and aging*

This grandmother hypothesis proposes that when food resources that were difficult for young juveniles to exploit became increasingly important in ancestral diets, mothers shared more of the food they found and processed with their young offspring. Help with provisioning then allowed earlier weaning and reduced birth spacing. Older females who were becoming infertile, and so not encumbered with infants themselves, could supply that help and have a large effect on their daughters' fertility. By this pathway selection would have favored increased allocation to physiological processes that buffer adults against mortality risk, slowing senescence through increases in somatic durability, maintenance, and repair (Kirkwood, 1977; Kirkwood and Holliday, 1979; Kirkwood and Rose, 1991).

This scenario assumes that australopithecine populations were in evolutionary equilibrium for the tradeoff between somatic maintenance and current reproduction. At this equilibrium, further reductions in the rates of senescence were too costly in current reproduction to be favored by selection. But then, the ecological changes that increased the importance of foods difficult for young juveniles to handle and increased the role of



maternal provisioning changed these pay-offs. Females who were slightly more vigorous as their own fertility was ending could increase the fertility of their daughters. Increased somatic effort would impose a cost on current reproductive effort: more vigorous longer-lived females would allocate less to their own current reproduction. But sufficient benefit from the help of elders could compensate that cost, resulting in selection for greater longevity. Net higher fertility for the child-bearers would give a selective advantage to lineages with increased vigor at later ages. This would slow further the already slow life history of our australopithecine ancestors.

Selection for greater longevity by this pathway could not delay the age-specific termination of fertility. Grandmothers with infants of their own would be less able to help their daughters. The cost of a more durable soma (lower reproductive effort from longer-lived mothers) in lineages with longer fertile spans would then go uncompensated. Thus, selection would favor less somatic effort and life histories collapse back to the ancestral chimpanzee-like equilibrium. Longer-lived lineages would maintain their slower life histories only by maintaining the ancestral age of menopause.

#### HUMAN/CHIMPANZEE COMPARISONS

This scenario depends on the inference that australopithecines had life histories similar to modern chimpanzees, while *H. erectus/ergaster* had life histories more similar to modern humans. Among the comparisons between humans and chimpanzees that are pertinent to it, I consider two here. If an ancestral age-specific fertility decline persisted through australopithecines to genus *Homo*, and if chimpanzees maintain a similar ancestral pattern, then we should be similar to chimpanzees in our age-related fertility declines. On the other hand, grandmothering in genus *Homo* is hypothesized to favor increased longevity. We should be different from chimpanzees in our somatic durability and age-specific somatic performance.

##### *Age-specific fertility decline*

Measurements of the day-specific probabilities of conception controlling intercourse behavior show fertility declines in women

beginning just after the mid-20s (Dunson et al., 2002). In industrial societies women in their 40s have few babies (e.g., Fretts et al., 1995), a pattern that also holds among modern foragers (Howell, 1979; Hill and Hurtado, 1996). Almost all last births are before 45. Data on chimpanzees are extremely limited by comparison. Conception rates per cycle in captivity are reported to be five times higher in females age 15–25 than in those age 35–48 (Graham, 1986). The ages of older adults are often not known in captivity, and known more rarely in the wild. Goodall (1986) estimated that Flo, the famous matriarch of the Gombe population, had her last birth in her early 40s. Flo's daughter Fifi, whose estimated birth is 1958, had her eighth baby in 1998. Sugiyama (1994) reports a few births to mothers estimated to be between 40 and 44 at Bossou. Based on those data, Gage (1998:209) estimates that "the end of the reproductive career... is similar for *Pan* and humans."

Comparing ages at menopause is more difficult, in part because menopause is harder to document even in humans. Treolar's (1981) dataset, based on longitudinal records, remains one of the best. Subjects initially recruited as students at the University of Minnesota recorded each menstruation and their daughters enrolled at menarche. In this well-nourished population, normal menopause ranged widely from early 40s to late 50s, with mean and mode for women not using hormone interventions about 51. Samples from other countries suggest there may be variation among different human populations. Some show earlier peaks and nutrition and other health practices may affect the timing of menopause, although measurement problems plague these comparisons (Gosden, 1985; Wood, 1994; Leidy, 1994).

In contrast to the large human datasets, there are few data on menopause in chimpanzees — in part because chimpanzees rarely live to menopausal ages. However, endocrine profiles published on two long captive subjects show them to be perimenopausal at ages 48 and 50, respectively (Gould et al., 1981), near the central tendencies of the human distribution. In addition to the two chimpanzees (*Pan troglodydes*), this study also included one bonobo (*Pan paniscus*). Estimated to be over 40 but younger than the other two, she was no longer cycling. Her hormone profiles and ovarian histology

confirmed that she had passed menopause. This provides another data point consistent with the premise that genus *Pan* and genus *Homo* have similar menopausal ages.

These limited data on age-specific fertility declines and menopause are consistent with the proposition that the ancestral age at menopause has been maintained in our lineage. However, some biochemical constraint, not an absence of directional selection, could also account for these similarities. Mammalian reproductive physiology makes this seem initially plausible. Although women, like most other mammals, reach menopause when a fixed stock of oocytes is depleted (Gosden, 1985; Richardson et al., 1987; vom Saal et al., 1994; Wood 1994), the similarity between human and chimpanzee menopause could result from an upper age limit on oocyte viability of about 50 years. But two kinds of evidence, interspecific variation in the size and depletion rates of initial oocyte stocks, and the longer fertile spans in other species suggest that selection can delay the age at menopause when there is net benefit for continued fertility at later ages.

First, the variation between and within species in both the size of the initial stock of oocytes and the rate at which these are lost is substantial (Finch, 1990; Finch and Kirkwood, 2000). Humans begin with an average of several million oocytes attained in the first few months of fetal life, reduced to less than a million before birth, with most of the remaining lost before puberty (Block, 1953). Number of ovulations is not a likely predictor of age at menopause, since less than 0.01% of the initial stock is ever actually ovulated. Across the mammals, the store of oocytes remaining at maturity varies with lifespan (Gosden and Telfer, 1987). Those stocks are larger in species with longer potential lives and the rate of loss during adulthood may be slower in longer-lived species (Gosden and Telfer, 1987). The cross-species variation suggests that selection can and does adjust both of these variables.

The second line of evidence consistent with the view that menopause at 50 is not set by physiological constraints on oocyte viability comes from other long-lived mammals. Both Asian (Sukumar et al., 1997) and African elephants (Moss, 2001) continue to have successful births into their seventh decade, 20 years longer than women do. In some cetaceans fertility continues to even later

ages. Antarctic fin whales are found pregnant into their 80s (Mizroch, 1981). With recent evidence that maximum lifespans reach well over 100 years in bowhead whales (George et al., 1999), data on late age fertility in this species will be of special interest.

Overall then, the available data on ages at last birth and menopause in chimpanzees show age-specific fertility declines in that species not substantially different from our own. Mammalian fertility, however, can extend to much older ages than it does in humans. This evidence is consistent with the argument that ancestral age-specific fertility declines have been maintained in our lineage, perhaps conserved by stabilizing selection.

#### *Distinctively human design for longevity*

While ages at terminal fertility and menopause appear to be similar in humans and chimpanzees, we differ from them and from other primates in our general vigor at these ages and our high probability of decades of menopausal survival (Pavelka and Fedigan, 1991). Goodall (1986:81) classified Gombe chimpanzees as moving from middle to old age at 33. Among the few in her study population who lived that long, she noted frailty, emaciation, slow movement, and difficulty in climbing. "Whether or not the lives of these individuals were complicated by disease during their last months, there can be little doubt that old age itself was primarily responsible for their deaths" (1986:104).

Wide recognition of human postmenopausal longevity has, however, not led to consensus that perimenopausal vigor and postcycling survival are distinctive characteristics of our evolutionary design. None dispute that we have "the maximum lifespan of the terrestrial animals" (Smith and Tompkins, 1995:258), but the reasons for this continue to be debated. I first review demographic data from various human populations to show that life expectancy at birth is a misleading index of longevity and that changes in mortality with age are regular enough to suggest a "characteristic pattern" in our species. Then, noting the possibility that a distinctively human pattern of adult mortality could be an artifact of our unusual economic interdependence and cooperation, I turn to physiological measures that index cross-species differences in the rate of senescence.

Demography. Some investigators argue that, rather than a species characteristic, human potential longevity is a recent novelty (e.g., Olshansky et al., 1998). In most of the US, western Europe, and Japan, newborns can now expect to live about 80 years, while historical demography and population profiles in traditional settings show life expectancies less than four decades. Life expectancies at birth, however, are strongly affected by rates of infant and juvenile mortality (e.g., Bailey, 1987; Smith DWE, 1993; Lee, 1997). There is no credible evidence of any change in maximum human lifespan over historical time (Austad, 1997).

French historical demography, for example, illustrates this point. Life expectancy was only 39 in 1850, compared to double that in 1985 (Keyfitz and Fleiger, 1968, 1990). The difference between these two time periods is not, however, a change in maximum life spans but a change in the rate of mortality at young ages. Since life expectancies are cohort averages, they reflect not only the long lives of those who die old but also the very short lives of those who die as infants and children. Even in mid-19th century France, with life expectancies less than 40, those who survived to adulthood had the prospect of a long life ahead. Of the girls who lived to adulthood, 72% then lived past 45, the age of terminal fertility, and the average number of years remaining for anyone who reached that age was more than two additional decades (Keyfitz and Fleiger, 1968, 1990).

Nineteenth-century France had an agrarian economy. If those vital rates depend on the use of domesticates, they would only have appeared within the last 10,000 years with the advent of agriculture. Demographic data from modern people who do not depend on domesticated food, however, show similar patterns of age-specific mortality. Three of the best-studied cases of modern hunter-gatherers, the !Kung of southern Africa (Howell, 1979), the Ache of South America (Hill and Hurtado, 1996), and the east African Hadza (Blurton Jones et al., 1992, 2002), represent populations in different environments with distinct recent genetic histories. All show patterns of age-specific survival very like those reported for 19th-century France. Life expectancies at birth are less than four decades in all these cases due to infant and juvenile mortality. On reaching adulthood most women live past their mid-40s, the age of terminal fertility,

and those who do have an *average* of more than 20 years of life still ahead.

Using Hamilton's (1966) method for computing the force of mortality, the instantaneous death rate for each age class, underscores the similarity in patterns of senescent mortality in human populations with diverse economies and recent genetic histories. Figure 1 plots demographic data from France in 1985, 1950, 1900, and 1850. Although life expectancies double over this time period, the age-specific mortality risks among adults are similar. Figure 2 shows the force of mortality calculated for the three hunter-gatherer populations, the !Kung (Howell, 1979), the Ache (Hill and Hurtado, 1996), and the Hadza (Blurton Jones et al., 2002).

The unusual postmenopausal longevity of humans is especially clear when we are compared with other primates (Pavelka and Fedigan, 1991, 1999). In macaque and baboon populations (Pavelka and Fedigan, 1999; Packer et al., 1998), reaching the age of terminal fertility is the luck of only a very few individuals. Less than 5% of those reaching adulthood live past that. Chimpanzees mature later and have longer adult lifespans than do macaques and baboons. In captivity a few more females live past childbearing age (Caro et al., 1995). But among chimpanzees in the wild (Hill et al., 2001) only about 5% of the adult females are past the likely age of terminal fertility — so few they are all but invisible statistically. This contrasts with about one-third of adult females beyond childbearing age in human foraging populations (Hawkes et al., 2003).

Figure 3 combines the hunter-gatherer cases, mid-19th century France, and the Taiwanese peasants of 1900, the dataset Hamilton (1966) used himself. While the differences among these modern human populations are not negligible, the similarities are striking. Similarities in the shape of changes in mortality with increasing age, even when the particular causes of death vary (Preston et al., 1972; Finch et al., 1990; Gavrilov and Gavrilov, 1991; Hill and Hurtado, 1996) seem a "species signature" of aging that is especially clear when the human populations are contrasted with chimpanzees (Fig. 3; chimpanzee data from Hill et al., 2001). Senescent mortality (the fraction of deaths above baseline — the lowest age-specific mortality) increases much earlier and more steeply for chimpanzees.




Fig. 1. Instantaneous age-specific mortality risk (Hamilton, 1966) calculated from historical demography (Keyfitz and Fleiger, 1968, 1990). Probability of death during a 5-year age class indicated on the y axis, age on the x axis.




Fig. 2. Instantaneous age-specific mortality risk (Hamilton, 1966) calculated for three contemporary hunter-gatherer populations. Probability of death during a 5-year age class indicated on the y axis, age on the x axis. !Kung from Howell (1979); Ache from Hill and Hurtado (1996); Hadza data Blurton Jones et al. (2002).

The inference from historical and ethnographic demography that substantial longevity is a characteristic feature of human populations has been challenged by age distributions of archaeological skeletal assemblages (Weiss, 1973; Trinkaus, 1986, 1995; Austad, 1997). Remains from individuals estimated to be over 60 at death are rare,

supporting skepticism about the generality of the demographic patterns found among living populations. But new sources of error are introduced in constructing population profiles from archaeological assemblages. Two especially important sources of bias are revealed by cases in which historical records provide independent evidence of




Fig. 3. Instantaneous age-specific mortality risk (Hamilton, 1966) for two preindustrial agricultural and three hunter-gatherer human populations and for wild chimpanzees. Probability of death during a 5-year age class indicated on the y axis, age on the x axis. French historical demograph from Keyfitz and Fleiger (1968); Taiwan peasant farmer data used by Hamilton (1966); !Kung from Howell (1979); Ache from Hill and Hurtado (1996); Hadza from Blurton Jones et al. (2002); chimpanzee data (smoothed) from Hill et al. (2001).

the age of individuals interred. Not only are the bones of the old and the young disproportionately unlikely to sustain long preservation, the ages of adults are also systematically underestimated (Walker et al., 1988; Paine, 1997). Standard aging techniques applied to samples of known ages illustrate the pervasiveness of the problem (Bocquet-Appel and Masset, 1982; Key et al., 1994). Analysts have repeatedly shown that the cemetery profiles do not represent sustainable living populations (e.g., Howell, 1982; Bermudez de Castro and Nicolas, 1997).

Biases in the other direction, novel features of the modern world that might extend longevity in ethnographically known foraging populations, have also been explored. Living people are now everywhere affected to some degree by global networks of interaction (Wolf, 1982; Schrire, 1994; Blurton Jones et al., 1996). Blurton Jones et al. (2002) examined the possible effects that interaction with agricultural neighbors as well as some access to Western medicines might have on Hadza demography. Even the most generous estimates of neighbor interactions, regional medical services, and the ethnographers' own possible effects

make only negligible differences in the population parameters initially reported (Blurton Jones et al., 2002).

These patterns are consistent with the argument that modern humans have a distinctive design for longevity. But human dependence on culture might provide an alternative explanation for potentially longer human lives. When Lee (1968) estimated that 10% of the people in his !Kung study population were over 60, he used the estimate to underline his observation that hunter-gatherers meet their needs more easily than often expected. He emphasized not only the time they spent in food procurement, but also their economic cooperation and food sharing. The longevity of humans might result from resource pooling and community support that reduces fatalities otherwise likely to follow illness, injury, or foraging failures. If greater human longevity results from distinctively human patterns of cooperation and interdependence, the demographic differences between foraging people and chimpanzees might greatly overestimate underlying patterns of age-specific frailty. A cultural safety net, rather than a fundamental shift in rates of senescence, could be the reason for our longer lives.

Physiological aging. If the difference between chimpanzee and human longevities reflects real differences in aging rates, and mechanisms of maintenance and repair are similar, differences in allocation to these mechanisms might be directly measurable. Evidence is mounting that “modulators of the rate of aging” are conserved over large evolutionary distances (Partridge and Gems, 2002:165). I summarize ideas and evidence about three physiological systems related to maintenance and repair: antioxidants, DNA repair, and age-related neuropathologies. All illustrate differences between humans and chimpanzees in somatic durability.

The accumulation of damage to cells as a result of metabolism is an important contributor to aging (Harman, 1957; Beckman and Ames, 1998; Finkel and Holbrook, 2000). This may explain how dietary restriction slows aging in a wide array of organisms (Holliday, 1989; Sohal and Weinruch, 1996; Yu, 1996; Lin et al., 2000; Hulbert and Else, 2000; Partridge and Gems, 2002). Cell lines from different mammalian species vary in their resistance to a variety of oxidative and nonoxidative stresses and the variation is correlated with species longevities (Kapahi et al., 1999). Both rates of free radical production (Barja et al., 1994) and the production of antioxidant mechanisms (Ku et al., 1994) also correlate with species longevity. This work provides a foundation for expecting design for greater longevity in humans than chimpanzees to be reflected in differences in the management of oxidative damage at the cellular level. Some data are available on antioxidant production among primates showing that levels of the circulating antioxidants plasma urate (Ames et al., 1981) and super oxide dismutase (Tolmasoff et al., 1989) are substantially higher in humans than in chimpanzees.

Damage to DNA accumulates with time, so variation in cell responses to this damage should be correlated with aging rates (Tice and Setlow, 1985). Investigators have used UV-irradiation to damage cell lines and then measured subsequent DNA repair. Cell lines from different mammalian species show differing amounts of repair depending on species longevities. In human cell lines DNA repair is much greater than in cell lines from other primates (Hart and Setlow, 1974). That sample did not include chimpanzees but it did include gorillas. To the extent that gorillas can serve as a proxy for chimpanzees (who

are themselves used here as a proxy for australopithecines), this is more evidence that humans are designed for greater longevity than were ancestral hominins.

Alzheimer-like neuropathologies increase with age in most primates just as they do in humans (Finch and Sapolsky, 1999). Unlike other primates, however, where the proliferation of senile plaques can begin before the age of menopause, these neuropathologies are generally delayed well past menopause in humans. When the same two chimpanzees found to be perimenopausal at ages 48 and 50 (Gould et al., 1981) died (at 56 and 59) they showed no sign of cognitive impairment (Gearing et al., 1994). However, autopsy revealed modest accumulation of brain amyloid (Gearing et al., 1994) which may represent presymptomatic or early symptomatic Alzheimer’s disease (Morris et al., 1996). This would be consistent with earlier aging in chimpanzees than humans. Finch and Sapolsky (1999) suggest that the delay in this kind of age-related decline in humans is associated with our unusually long postmenopausal survival linked to the benefits of grandmothing. They also hypothesize that the several genetic variants of apolipoprotein E that are found in humans but not in other primates may be implicated in the protection against these age-related neuropathologies in our species. One of the apo E variants most like the single isoform found in other primates is associated with an array of disease susceptibilities that are consistent with this hypothesis (Finch and Sapolsky, 1999).

## CONCLUSIONS

Like Washburn’s elaboration of the hunting hypothesis, a grandmother hypothesis assumes that changing ecological circumstances favored a shift in foraging strategies that gave rise to our genus, with assistance for juveniles from kin other than mothers driving the shift. But this grandmother hypothesis uses a demographic perspective on life-history variation that gives our low adult mortality rates and remarkable longevity special importance.

The claim is that members of our genus began living longer *not* because they were kept alive when frail — although this does now happen, increasingly in many current populations. Instead, a change in selection pressures that favored increased

allocation to somatic durability resulted in slower aging in our genus. Greater longevity in turn favored later ages of maturity. Australopithecines with chimpanzee-like life histories had maintained a different equilibrium in which reproductive and other physiological systems senesced in tandem. When ecological changes resulted in more mother-child food sharing, the cost of variants that reduced senescent mortality in ancestral populations were outweighed by the benefits due to more vigorous grandmothers' help. This resulted in the evolution of genus *Homo*.

Instead of linking late maturity to the developmental requirements of large-brained juveniles, the grandmother hypothesis derives our late maturity from our design for longevity. Later maturity also means that older juveniles, gaining benefits from delaying their own reproduction, could provide another source of help for younger siblings. Fathers may have helped sometimes. But ancient men, like other males (Hawkes et al., 1995), especially other primates (Smuts, 1987, Smuts and Gubernick, 1992; Van Schaik and Paul, 1996), and like modern men (Hurtado and Hill, 1992; Bliege Bird, 1999; Blurton Jones et al., 2000; Hawkes et al., 2001), could not have escaped the pressures of male competition that often make mating effort take priority over paternal investment (Williams, 1966a; Trivers, 1972; Hawkes and Bliege Bird, 2002; O'Connell et al., 2002b). The grandmother hypothesis, unlike the hunting hypothesis, does not assume that cooperation in child-rearing either depends on or necessarily gives rise to nuclear families.

Hrdy (1999, 2001) has characterized humans as cooperative breeders. Noting that nonmaternal helpers, or allomothers, play an especially important role in rearing human children, she links a greater human ambivalence about maternal investment in particular offspring to our greater need for help. "A human mother's commitment to her infant should be linked to how much social support she herself can expect" (Hrdy, 2001:58). Most other primates postpone the next offspring as long as the previous one is still dependent. The grandmother hypothesis explains how we came to do otherwise, deriving the evolution of our need for help from the postmenopausal longevity that supplied more candidates for allomothering. Vigorous peri- and postmenopausal women, overlapping infants

and toddlers, as well as prereproductive adolescents, are part of a systematically inter-related suite of life-history features. Together these features make social circumstances especially important, and so make maternal sensitivity to them especially advantageous.

A growing proportion of adults in senior age ranks in many contemporary human populations is clearly a recent novelty (Vaupel et al., 1998; Oeppen and Vaupel 2002). This large fraction of elders now surviving in spite of senescent frailty presents economic, medical, political, and social challenges that can hardly be overestimated (Tuljapurkar et al., 2000). But those facts should not obscure the strength of the evidence that adult lives substantially longer than those of the other apes are "normal" for our species, yet our age-specific fertility decline and menopause appears to differ little from chimpanzees. The models and evidence reviewed here employ the widely used assumption that chimpanzees can serve as a life-history analog to human ancestors. If australopithecines had ape-like life histories, the effects of senescence on adult mortalities could have been very large.

Demographic comparisons between foraging chimpanzees and people (Fig. 3) show not only the earlier, steeper climb in senescent mortality in chimpanzees, the baseline "initial mortality" rate is also higher for chimpanzees (Hill et al., 2001). While this lowest rate is conventionally used to estimate extrinsic mortality risk (e.g., Ricklefs, 1998), disposable soma ideas suggest that it would also include an intrinsic component in long-lived species. If mechanisms of maintenance and repair develop early in ontogeny, then more effective mechanisms for buffering mortality risks in adulthood would likely reduce the initial mortality rate as well (Finch et al., 1990).

Washburn argued effectively for the use of comparisons with other primates to discover what happened in human evolution. When he advanced the hunting hypothesis the frequency and importance of hunting among chimpanzees, now so well documented (e.g., Goodall, 1986; Boesch and Boesch, 1989; Stanford, 1996, 1999; Mitani et al., 2002), was unknown. In addition to the essential value of those and other behavioral comparisons between these species (e.g., de Waal 2001), genetic comparisons now promise the possibility of detailed evidence about human evolution (Varki, 2000; Gagneux and Varki,

2001; Enard et al., 2002). More work focused especially on life-history comparisons is also in order, including comparisons in age-related physiology (Erwin et al., 2001) and in molecular and cellular processes associated with somatic development, maintenance, and repair.

#### ACKNOWLEDGMENTS

I thank H.P. Alvarez, N.G. Blurton Jones, E.L. Charnov, C.E. Finch, N. Howell, T.B.L. Kirkwood, R.G. Klein, C.W. Kuzawa, L.G. Moore, J.F. O'Connell, and G.C. Williams for useful comments and good advice.

#### LITERATURE CITED

- Aiello LC, Wells JCK. 2002. Energetics and the evolution of genus *Homo*. *Annu Rev Anthropol* 31:323–338.
- Aiello LC, Wheeler PK. 1995. The expensive-tissue hypothesis: the brain and the digestive system in human and primate evolution. *Curr Anthropol* 36:199–221.
- Alvarez HP. 2000. Grandmother hypothesis and primate life histories. *Am J Phys Anthropol* 133:435–450.
- Ames BN, Cathcart R, Schwiers E, Hochstein P. 1981. Uric acid provides an antioxidant defense in humans against oxidant- and radical-caused aging and cancer: a hypothesis. *Proc Natl Acad Sci USA* 78:6858–6862.
- Anton SC. 2001. Growth and life history in *Homo erectus*. *AAPA Abstracts Am J Phys Anthropol* 114:32–33.
- Austad SN. 1997. Postreproductive survival. In: Wachter KW, Finch CE, editors. *Between Zeus and the salmon*. Washington, DC: National Academy Press. p 161–174.
- Bailey WG. 1987. Human longevity from antiquity to the modern lab: a selected, annotated bibliography. New York: Greenwood Press.
- Barja G, Cadenas S, Rojas C, Perez-Campo R, Lopez-Torres M. 1994. Low mitochondrial free radical production per unit O<sub>2</sub> consumption can explain the simultaneous presence of high longevity and high aerobic metabolic rate in birds. *Free Rad Res* 21:317–327.
- Beckman, KB, Ames BN. 1998. The free radical theory of aging matures. *Phys Rev* 78:547–581.
- Bermudez de Castro, JM, Nicolas, ME. 1997. Paleodemography of the Atapuerca-Sima de los Huesos Middle Pleistocene hominid sample. *J Hum Evol* 33:333–355.
- Bliege Bird R. 1999. Cooperation and conflict: the behavioral ecology of the sexual division of labor. *Evol Anthropol* 8:65–75.
- Blumenshine RJ. 1987. Characteristics of an early hominid scavenging niche. *Curr Anthropol* 28:383–407.
- Blurton Jones NG, Hawkes K, O'Connell JF. 1989. Studying costs of children in two foraging societies: implications for schedules of reproduction. In: Standon V, Foley R, editors. *Comparative socioecology of mammals and man*. London: Blackwell. p 365–390.
- Blurton Jones NG, Smith LC, O'Connell JF, Hawkes K, Kamazura C. 1992. Demography of the Hadza, an increasing and high density population of savanna foragers. *Am J Phys Anthropol* 89:159–181.
- Blurton Jones NG, Hawkes K, O'Connell JF. 1996. The global process and local ecology: how should we explain differences between the Hadza and !Kung? In: Kent S, editor. *Cultural diversity among twentieth century foragers: an African perspective*. Cambridge, UK: Cambridge University Press. p 159–187.
- Blurton Jones NG, Marlowe F, Hawkes K, O'Connell JF. 2000. Paternal investment and hunter-gatherer divorce. In: Cronk L, Chagnon N, Irons W, editors. *Adaptation and human behavior; an anthropological perspective*. New York: Aldine de Gruyter. p 61–90.
- Blurton Jones NG, Hawkes K, O'Connell JF. 2002. The antiquity of post-reproductive life: are there modern impacts on hunter-gatherer post-reproductive life spans? *Am J Hum Biol* 14:184–205.
- Binford LR. 1981. *Bones: ancient men and modern myths*. New York: Academic Press.
- Block E. 1953. Quantitative morphological investigation of the follicular system in newborn female infants. *Acta Anat* 17:201–206.
- Blueweiss L, Fox H, Kudza V, Nakashima D, Peters R, Sams S. 1978. Relationships between body size and some life history parameters. *Oecologia* 37:257–272.
- Bocquet-Appel J-P, Masset C. 1982. Farewell to paleodemography. *J Hum Evol* 11:321–333.
- Boesch C, Boesch H. 1989. Hunting behavior of wild chimpanzees in Tai National Park. *Am J Phys Anthropol* 78:547–573.
- Bogin B. 1999. *Patterns of human growth*, 2nd edition. Cambridge, UK: Cambridge University Press.
- Bogin B. 2001. *The growth of humanity*. New York: Wiley-Liss.
- Bonner JT. 1965. *Size and cycle: an essay in the structure of biology*. Princeton, NJ: Princeton University Press.
- Bromage T. 1987. The biological and chronological maturation of early hominids. *J Hum Evol* 16:257–272.
- Bromage T, Dean C. 1985. Reevaluation of the age at death of Plio-Pleistocene fossil hominids. *Nature* 317:525–538.
- Carey JR, Judge DS. 2001. *Principles of biodemography with special reference to human longevity*. Population; An English selection 13:9–24.
- Caro TM, Sellen DW, Parish A, Frank R, Brown DM, Voland E, Borgerhoff Mulder M. 1995. Termination of reproduction in nonhuman and human female primates. *Int J Primatol* 16:205–220.
- Cartmill M. 1993. *A view to a death in the morning: hunting and nature through history*. Cambridge, MA: Harvard University Press.
- Charnov EL. 1991. Evolution of life history variation in female mammals. *Proc Natl Acad Sci USA* 88:1134–1137.
- Charnov EL. 1993. Life history invariants: some explorations of symmetry in evolutionary ecology. Oxford: Oxford University Press.
- Charnov EL. 1997. Trade-off-invariant rules for evolutionarily stable life histories. *Nature* 387:393–394.
- Charnov EL. 2001. Evolution of mammal life histories. *Evol Ecol Res* 3:521–535.
- Charnov EL, Berrigan D. 1991. Dimensionless numbers and the assembly rules for life histories. *Philos Trans R Soc Lond B* 332:241–248.
- Charnov EL, Berrigan D. 1993. Why do female primates have such long lifespans and so few babies? Or, life in the slow lane. *Evol Anthropol* 1:191–194.
- Charnov EL, Schaffer WM. 1973. Life history consequences of natural selection: Cole's result revisited. *Am Nat* 107:791–793.
- Clegg M, Aiello LC. 1999. A comparison of the Nariokotome *Homo erectus* with juveniles from a modern human population. *Am J Phys Anthropol* 110:81–93.



- Cole LC. 1954. The population consequences of life history phenomena. *Q Rev Biol* 29:103-137.
- Collard M, Wood B. 1999. Grades among the African early hominids. In: Bromage TG, Friedemann S, editors. African biogeography, climate change, and human evolution. Oxford: Oxford University Press. p 316-327.
- Dart R. 1949. The predatory implemental technique of *Australopithecus*. *Am J Phys Anthropol* 7:1-38.
- Dart R. 1953. The predatory transition from ape to man. *Int Anthropol Linguist Rev* 1:201-217.
- De Waal FBM (ed.). 2001. Tree of origin: what primate behavior can tell us about human social evolution. Cambridge, MA: Harvard University Press.
- Dean C, Leakey MG, Reid D, Schrenk F, Schwartz GT, Stringer C, Walker A. 2001. Growth processes in teeth distinguish modern humans from *Homo erectus* and earlier hominins. *Nature* 414:628-631.
- Dominguez-Rodrigo M. 2002. Hunting and scavenging by early humans: the state of the debate. *J World Prehist* 16:1-54.
- Dunson DB, Colombo B, Baird DD. 2002. Changes with age in the level and duration of fertility in the menstrual cycle. *Hum Reprod* 17:1399-1403.
- Eisenberg JF. 1981. The mammalian radiations. Chicago: University of Chicago Press.
- Enard W, Przeworski M, Fisher SE, Lai CSL, Wiebe V, Kitano T, Monaco AP, Paabo S. 2002. Molecular evolution of FOXP2, a gene involved in speech and language. *Nature* 418:469-472.
- Erwin JM, Hof PR, Ely JJ, Perl DP. 2002. One gerontology: advancing understanding of aging through studies of great apes and other primates. In: Erwin JM, Hof PR, editors. Aging in nonhuman primates. Interdisciplinary topics in gerontology. Basel: Karger. 31:1-21.
- Finch CE. 1990. Longevity, senescence, and the genome. Chicago: University of Chicago Press.
- Finch CE. 1997. Comparative perspectives on plasticity in human aging and life spans. In: Wachter KW, Finch CE, editors. Between Zeus and the salmon. Washington, DC: National Academy Press. p 245-268.
- Finch CE, Kirkwood TBL. 2000. Chance, development, and aging. New York: Oxford University Press.
- Finch CE, Sapolsky R. 1999. The evolution of Alzheimer disease, the reproductive schedule, and apoE isoforms. *Neurobiol Aging* 20:407-428.
- Finch CE, Tanzi R. 1997. Genetics of aging. *Science* 278:407-411.
- Finch CE, Pike MC, Whitten M. 1990. Slow mortality rate accelerations during aging in some animals approximate that of humans. *Science* 249:902-905.
- Finkel T, Holbrook NJ. 2000. Oxidants, oxidative stress and the biology of ageing. *Nature* 408:239-246.
- Fretts RC, Schmittiel J, McLean FH, Usher RH, Goldman MB. 1995. Increased maternal age and the risk of fetal death. *N Engl J Med* 333:953-957.
- Gabunia L, Vekua A, Lordkipanidze D, Swisher CC III, Ferring R, Justus A, Nioradze M, Tvalchrelidze M, Antón SC, Bosinski G, Jöris O, de Lumley M-A, Majsuradze G, Mouskhelishvili A. 2000. Earliest Pleistocene hominid cranial remains from Dmanisi, Republic of Georgia: taxonomy, geological setting, and age. *Science* 288:1019-1025.
- Gadgil M, Bosser W. 1970. Life history consequences of natural selection. *Am Nat* 104:1-24.
- Gage TB. 1998. The comparative demography of primates: with some comments on the evolution of life histories. *Annu Rev Anthropol* 27:197-221.
- Gagneux P, Varki A. 2001. Genetic differences between humans and great apes. *Mol Phylogenet Evol* 18:2-13.
- Gavrillo LA, Gavrillo NS. 1991. The biology of lifespan: a quantitative approach. Amsterdam: Harwood Academic.
- Gearing M, Rebeck WG, Hyman BT, Tigges J, Mirra SS. 1994. Neuropathology and apolipoprotein E profile of aged chimpanzees: implications for Alzheimer disease. *Proc Natl Acad Sci USA* 91:9382-9386.
- George JC, Bada J, Zeh J, Scott L, Brown L, O'Hara T, Suydam R. 1999. Age and growth estimates of bowhead whales (*Balaena mysticetus*) via aspartic acid racemization. *Can J Zool* 77:571-580.
- Goodall J. 1986. The chimpanzees of Gombe. Cambridge, MA: Harvard University Press.
- Gosden RG. 1985. Biology of menopause: the causes and consequences of ovarian ageing. London: Academic Press.
- Gosden RG. 1996. Cheating time: science, sex and ageing. London: Macmillan.
- Gosden RG, Telfer E. 1987. Numbers of follicles and oocytes in mammalian ovaries and their allometric relationships. *J Zool Lond* 211:169-175.
- Gould SJ. 1977. Ontogeny and phylogeny. Cambridge: Belknap Press of Harvard.
- Gould KG, Flint M, Graham, CE. 1981. Chimpanzee reproductive senescence: A possible model for the evolution of the menopause. *Maturitas* 3:157-166.
- Gowaty P. 1996. Battles of the sexes and the origins of monogamy. In: Black JM, editor. Partnerships in birds. Oxford: Oxford University Press. p 21-52.
- Graham CE. 1986. Endocrinology of reproductive senescence. In: Dukelow WR, Erwin J, editors. Comparative primate biology, vol. 3. Reproduction and development. New York: AR Liss. p 93-99.
- Haig D. 1993. Genetic conflicts of human pregnancy. *Q Rev Biol* 68:495-532.
- Haile-Selassie Y. 2001. Late Miocene hominids from the middle Awash, Ethiopia. *Nature* 412:176-181.
- Hamilton WD. 1966. The molding of senescence by natural selection. *J Theor Biol* 12:12-45.
- Harman D. 1957. Aging: a theory based on free radical and radiation chemistry. *J Gerontol* 2:298-300.
- Hart RW, Setlow RB. 1974. Correlation between deoxyribonucleic acid excision repair and lifespan in a number of mammalian species. *Proc Natl Acad Sci USA* 71:2169-2173.
- Harvey PH, Clutton-Brock TH. 1985. Life history variation in primates. *Evolution* 39:559-581.
- Harvey PH, Read AF. 1988. How and why do mammalian life histories vary? In: Boyce MS, editor. Evolution of life histories: patterns and process from mammals. New Haven, CT: Yale University Press. p 213-232.
- Harvey PH, Purvis A. 1999. Understanding the ecological and evolutionary reasons for life history variation: mammals as a case study. In: McGlade J, editor. Advanced ecological theory: principles and applications. Oxford: Blackwell Science. p 232-248.
- Harvey PH, Zammuto RM. 1985. Patterns of mortality and age at first reproduction in natural populations of mammals. *Nature* 315:319-320.
- Harvey PH, Promislow DEL, Read AF. 1989. Causes and correlates of life history differences in mammals. In: Standon V, Foley R, editors. Comparative socioecology of mammals and man. London: Blackwell. p 305-318.
- Hawkes K, Bliege Bird R. 2002. Showing-off, handicap signaling, and the evolution of men's work. *Evol Anthropol* 11:58-67.
- Hawkes K, O'Connell JF, Blurton Jones NG. 1989. Hardworking Hadza grandmothers. In: Standon V, Foley R, editors. Comparative socioecology of mammals and man. London: Blackwell. p 341-366.

- Hawkes K, O'Connell JF, Blurton Jones NG. 1995. Hadza children's foraging: juvenile dependency, social arrangements and mobility among hunter-gatherers. *Curr Anthropol* 36:688-700.
- Hawkes K, O'Connell JF, Blurton Jones NG. 1997. Hadza women's time allocation, offspring provisioning, and the evolution of post-menopausal lifespans. *Curr Anthropol* 38:551-578.
- Hawkes K, O'Connell JF, Blurton Jones NG, Charnov EL, Alvarez HP. 1998. Grandmothering, menopause, and the evolution of human life histories. *Proc Natl Acad Sci USA* 95:1336-1339.
- Hawkes K, O'Connell JF, Blurton Jones NG. 2001. Hunting and nuclear families: some lessons from the Hadza about men's work. *Curr Anthropol* 42:681-709.
- Hawkes K, O'Connell JF, Blurton Jones NG. 2003. The evolution of human life histories: primate tradeoffs, grandmothering socioecology, and the fossil record. In: Kappeler P, Pereira M, editors. *Primate life histories and socioecology*. Chicago: University of Chicago Press. p 204-227.
- Herskind AM, McGue M, Holm NV, Sorensen TIA, Harvald B, Vaupel JW. 1996. The heritability of human longevity: A population based study of 2,872 Danish twin pairs born 1870-1900. *Hum Genet* 97:319-323.
- Hill K. 1982. Hunting and human evolution. *J Hum Evol* 11:521-544.
- Hill K, Hurtado AM. 1991. The evolution of premature reproductive senescence and menopause in human females: an evaluation of the "grandmother" hypothesis. *Hum Nat* 2:313-350.
- Hill K, Hurtado AM. 1996. Ache life history: the ecology and demography of a foraging people. Hawthorne, NY: Aldine de Gruyter.
- Hill K, Hurtado AM. 1999. Packer and colleagues' model of menopause for humans. *Hum Nat* 10:199-204.
- Hill K, Boesch C, Goodall J, Pusey A, Williams J, Wrangham R. 2001. Mortality rates among wild chimpanzees. *J Hum Evol* 40:437-450.
- Holliday R. 1989. Food, reproduction and longevity; is the extended lifespan of calorie-restricted animals an evolutionary adaptation? *BioEssays* 10:125-172.
- Holliday R. 1995. *Understanding Ageing*. Cambridge, UK: Cambridge University Press.
- Howell N. 1979. *Demography of the Dobe !Kung*. New York: Academic Press.
- Howell N. 1982. Village composition implied by a paleodemographic life table: the Libben Site. *Am J Phys Anthropol* 59, 263-269.
- Hrdy SB. 1981. *The woman that never evolved*. Cambridge, MA: Harvard University Press.
- Hrdy SB. 1999. *Mother Nature: a history of mothers, infants, and natural selection*. New York: Pantheon.
- Hrdy SB. 2001. Mothers and others. *Nat Hist* 110: 50-64.
- Hulbert AJ, Else PL. 2000. Mechanisms underlying the cost of living in animals. *Annu Rev Physiol* 62: 207-235.
- Hurtado AM, Hill K. 1992. Paternal effect on offspring survivorship among the Ache and Hiwi hunter-gatherers: implications for modeling pair-bond stability. In: Hewlett B, editor. *Father-child relations: cultural and biosocial contexts*. New York: Aldine de Gruyter. p 31-55.
- Isaac G L. 1978. The food sharing behavior of proto-human hominids. *Sci Am* 238, 4:90-108.
- Isaac G L. 1984. The archaeology of human origins: studies of the Lower Pleistocene in East Africa, 1971-1981. *Adv World Archaeol* 3:1-87.
- Jamison CS, Cornell LL, Jamison PL, Nakazato H. 2002. Are all grandmothers equal? A review and a preliminary test of the "grandmother hypothesis in Tokugawa Japan. *Am J Phys Anthropol* 119:67-76.
- Johanson DC, White TD. 1979. A systematic assessment of early African hominids. *Science* 202:321-330.
- Judge DS, Carey JR. 2000. Postreproductive life predicted by primate patterns. *J Gerontol Biol Sci* 55A:B201-B209.
- Kapahi P, Boulton ME, Kirkwood TBL. 1999. Positive correlation between mammalian life span and cellular resistance to stress. *Free Rad Biol Med* 26:495-500.
- Kaplan H. 1997. The evolution of the human life course. In: Wachter KW, Finch CE, editors. *Between Zeus and the salmon*. Washington, DC: National Academy Press. p 175-211.
- Kaplan H, Hill K, Lancaster J, Hurtado AM. 2000. A theory of human life history evolution: diet, intelligence, and longevity. *Evol Anthropol* 9:156-185.
- Key CA, Aiello L, Molleson T. 1994. Cranial suture closure and its implications for age estimation. *Int J Osteoarchaeol* 4:193-207.
- Keyfitz N, Fleiger W. 1968. *World population: an analysis of vital data*. Chicago: University of Chicago Press.
- Keyfitz N, Fleiger W. 1990. *World population growth and aging: demographic trends in the late twentieth century*. Chicago: University of Chicago Press.
- Kirkwood TBL. 1977. Evolution of aging. *Nature* 270:301-302.
- Kirkwood TBL. 1981. Repair and its evolution: survival versus reproduction. In: Townsend CR, Calow P, editors. *Physiological ecology: an evolutionary approach to resource use*. Oxford: Blackwell Scientific. p 165-189.
- Kirkwood, TBL, Holliday R. 1979. The evolution of aging and longevity. *Proc R Soc Lond B* 205:532-546.
- Kirkwood, TBL, Rose MR. 1991. Evolution of senescence: late survival sacrificed for reproduction. *Philos Trans R Soc Lond B* 332:15-24.
- Klein RG. 1999. *The human career: human biological and cultural origins*, 2nd ed. Chicago: University of Chicago Press.
- Klein RG. 2000. Archaeology and the evolution of human behavior. *Evol Anthropol* 9:17-36.
- Kozlowski J, Weiner J. 1997. Interspecific allometries are byproducts of body size optimization. *Am Nat* 149:352-380.
- Ku H-H, Brunk TU, Sohal RS. 1993. Relationship between mitochondrial superoxide and hydrogen peroxide production and longevity of mammalian species. *Free Rad Biol Med* 15:621-627.
- Lancaster JB, Lancaster C. 1983. Parental investment: the hominid adaptation. In: Ortner DJ, editor. *How humans adapt: biocultural odyssey*. Washington, DC: Smithsonian Institution Press. p 33-56.
- Lancaster JB, Lancaster C. 1987. The watershed: change in parental-investment and family formation strategies in the course of human evolution. In: Lancaster JB, Altmann J, Rossi AS, Sherrod LR, editors. *Parenting across the life span: biosocial dimensions*. Hawthorne, NY: Aldine de Gruyter. p 187-205.
- Larick R, Ciochon R, Zaim Y, Sudijono, Suminto, Rizal Y, Aziz F, Reagan M, Heizler M. 2001. Early Pleistocene 40Ar/39Ar ages for Babang Formation hominins, Central Java, Indonesia. *Proc Natl Acad Sci USA* 98:4866-4871.
- Lee RB. 1968. What hunters do for a living, or how to make out on scarce resources. In: Lee RB, DeVore I, editors. *Man the hunter*. Chicago: Aldine. p 30-48.
- Lee RD. 1997. Intergeneration relations and the elderly. In: Wachter KW, Finch CE, editors. *Between Zeus and the salmon*. Washington, DC: National Academy Press. p 212-233.

- Lee-Thorp J. 2002. Hominid dietary niches from proxy chemical indicators in fossils: the Swartkrans example. In: Ungar PS, Teaford MF, editors. Human diet: its origin and evolution. Westport, CT: Bergin & Garvey. p 123–142.
- Lee-Thorp J, Thackeray JF, van der Merwe N. 2000. The hunters and the hunted revisited. *J Hum Evol* 30:565–576.
- Leidy LE. 1994. Biological aspects of menopause: across the lifespan. *Annu Rev Anthropol* 23:231–253.
- Leigh SR. 2001. Evolution of human growth. *Evol Anthropol* 10:223–236.
- Lieberman DE. 2002. Whither the evolution of growth and development? *Evol Anthropol* 11:246–248.
- Lin SJ, Defossez PA, Guarente L. 2000. Requirement of NAD and SIR2 for lifespan extension by calorie restriction in *Saccharomyces cerevisiae*. *Science* 289:2126–2128.
- Mesnick SL. 1997. Sexual alliances: evidence and evolutionary implications. In: Gowaty PA, editor. *Feminism and evolutionary biology: boundaries, intersections, and frontiers*. New York: Chapman Hall. p 207–257.
- McArthur RH, Wilson EO. 1967. *Theory of island biogeography*. Princeton, NJ: Princeton University Press.
- McHenry HM. 1992. Body size and proportions in early hominids. *Am J Phys Anthropol* 87:407–431.
- McHenry HM. 1994. Behavioral ecological implications of early hominid body size. *J Hum Evol* 27:77–87.
- Milton K. 1999. A hypothesis to explain the role of meat-eating in human evolution. *Evol Anthropol* 8:11–21.
- Mitani JC, Watts DP, Muller MN. 2002. Recent developments in the study of wild chimpanzee behavior. *Evol Anthropol* 11:9–25.
- Mizroch SA. 1981. Analysis of some biological parameters in the Antarctic fin whale. Report of the International Whaling Commission 31:425–434.
- Morris JC, Storandt M, McKeel DW Jr, Rubin EH, Price JL, Brant EA, Berg L. 1996. Cerebral amyloid deposition and diffuse plaques in “normal” aging: evidence for presymptomatic and very mild Alzheimer’s disease. *Neurology* 46:707–719.
- Moss CJ. 2001. The demography of an African elephant (*Loxodonta africana*) population in Amboseli, Kenya. *J Zool* 255:145–156.
- O’Connell JF, Hawkes K, Blurton Jones NG. 1999. Grandmothering and the evolution of *Homo erectus*. *J Hum Evol* 36:461–485.
- O’Connell JF, Hawkes K, Blurton Jones NG. 2002a. Meat eating, grandmothering and the evolution of early human diets. In: Ungar PS, Teaford MF, editors. *Human diet: its origin and evolution*. Westport, CT: Bergin & Garvey. p 49–60.
- O’Connell JF, Hawkes K, Lupo KD, Blurton Jones NG. 2002b. Male strategies and Plio-Pleistocene archaeology. *J Hum Evol* 43:831–872.
- Oeppen J, Vaupel JW. 2002. Broken limits to life expectancy. *Science* 296:1029–1031.
- Olshansky SJ, Carnes BA, Grahn D. 1998. Confronting the boundaries of human longevity. *Am Sci* 86:52–61.
- Packer C, Tatar M, Collins A. 1998. Reproductive cessation in female mammals. *Nature* 392:807–811.
- Paine RR (ed.). 1997. *Integrating archaeological demography: multidisciplinary approaches to prehistoric population*. Occasional Paper 24, Center for Archaeological Investigations. Carbondale: Southern Illinois University.
- Palombi RA. 1999. Infanticide and the evolution of pair bonds in nonhuman primates. *Evol Anthropol* 7:117–129.
- Partridge L, Gems D. 2002. Mechanisms of ageing: public or private? *Nat Rev Genet* 3:165–175.
- Pavelka MSM, Fedigan LM. 1991. Menopause: a comparative lifehistory perspective. *Yearb Phys Anthropol* 34:13–38.
- Pavelka MSM, Fedigan LM. 1999. Reproductive termination in female Japanese monkeys: a comparative life history perspective. *Am J Phys Anthropol* 109:455–464.
- Peccei JS. 2001. Menopause: adaptation of epiphenomenon? *Evol Anthropol* 10:43–57.
- Pianka ER. 1970. On “r” and “K” selection. *Am Nat* 104:592–597.
- Preston S, Keyfitz N, Schoen R. 1972. *Causes of death: life tables for national populations*. New York: Academic Press.
- Promislow DEL, Read A, Harvey PH. 1989. Life history variation in placental mammals: unifying the data with theory. In: Harvey PH, Partridge L, editors. *Oxford surveys in evolutionary biology*. New York: Oxford University Press. 6:13–31.
- Purvis A, Harvey PH. 1995. Mammal life-history evolution: a comparative test of Charnov’s model. *J Zool Lond* 237:259–283.
- Read AF, Harvey PH. 1989. Life history differences among the eutherian radiations. *J Zool Lond* 219:329–353.
- Richardson SJ, Senikas V, Nelson JF. 1987. Follicular depletion during the menopausal transition: evidence for accelerated loss and ultimate exhaustion. *J Clin Endocrinol Metab* 65:1231–1237.
- Ricklefs RE. 1998. Evolutionary theories of aging: confirmation of a fundamental prediction, with implications for the genetic basis and evolution of life span. *Am Nat* 152:24–44.
- Rightmire CP. 1990. *The evolution of Homo erectus: comparative anatomical studies of an extinct species*. Cambridge, UK: Cambridge University Press.
- Rogers AR. 1993. Why menopause? *Evol Ecol* 7:406–420.
- Rose MR. 1991. *Evolutionary biology of aging*. New York: Oxford University Press.
- Ruff CB, Trinkhaus E, Holliday TW. 1997. Body mass and encephalization in Pleistocene Homo. *Nature* 387:173–176.
- Saether B-E. 1988. Pattern of co-variation between life-history traits of European birds. *Nature* 331:616–617.
- Schrire C (ed.). 1984. *Past and present in hunter gatherer studies*. Orlando: Academic Press.
- Sear R, Mace R, McGregor IA. 2000. Maternal grandmothers improve the nutritional status of children in rural Gambia. *Proc R Soc Lond B* 267:1641–1647.
- Senut B, Pickford M, Gommery D, Mein P, Cheboi K, Coppens Y. 2001. First hominid from the Miocene (Lukeino Formation, Kenya). *Comptes Rendus de l’Academie des Sciences, Series IIA — Earth and Planetary Science* 332. 2:137–144.
- Service PM, Hutchinson EW, McKinley MD, Rose MR. 1985. Resistance to environmental stress in *Drosophila melanogaster* selected for postponed senescence. *Physiol Zool* 58:380–389.
- Shanley DP, Kirkwood TBL. 2001. Evolution of the human menopause. *BioEssays* 23:282–287.
- Shipman P. 1986. Scavenging or hunting in early hominids: Theoretical framework and tests. *Am Anthropol* 88:27–43.
- Shipman P, Walker A. 1989. The costs of becoming a predator. *J Hum Evol* 18:373–392.
- Smith BH. 1986. Denatal development in *Australopithecus* and early *Homo*. *Nature* 323:327–330.
- Smith BH. 1987. Dental development in fossil hominids. *Am J Phys Anthropol* 72:255.
- Smith BH. 1989. Dental development as a measure of life history in primates. *Evolution* 43:683–688.
- Smith BH. 1991. Dental development and the evolution of life history in Hominidae. *Am J Phys Anthropol* 86:157–174.

- Smith BH. 1993. The physiological age of KNM-15000. In: Walker A, Leakey R, editors. *The Nariokotome Homo erectus* skeleton. Cambridge, MA: Harvard University Press. p 195-220.
- Smith DWE. 1993. Human longevity. New York: Oxford University Press.
- Smith BH. 1994. The sequence of eruption of teeth in *Macaca*, *Pan*, *Australopithecus*, and *Homo*: its evolutionary significance. *Am J Hum Biol* 6:61-76.
- Smith BH, Tompkins RL. 1995. Toward a life history of the hominidae. *Annu Rev Anthropol* 24:257-279.
- Smuts BB. 1985. Sex and friendship in baboons. New York: Aldine.
- Smuts BB. 1987. Sexual competition and mate choice. In: Smuts BB, Cheney DL, Seyfarth R, Wrangham RW, Struthsaker TT, editors. *Primate societies*. Chicago: University of Chicago Press. p 385-399.
- Smuts BB. 1992. Male aggression against women: an evolutionary perspective. *Hum Nat* 3:1-44.
- Smuts BB, Gubernick D. 1992. Male-infant relationships in nonhuman primates: Paternal investment or mating effort? In: Hewlett B, editor. *Father-child relations: cultural and biosocial contexts*. New York: Aldine de Gruyter. p 1-30.
- Sohol RS, Weinruch R. 1996. Oxidative stress, caloric restriction and aging. *Science* 273:59-63.
- Stanford C. 1996. The hunting ecology of wild chimpanzees: implications for the evolutionary ecology of Pliocene hominids. *Am Anthropol* 98:96-113.
- Stanford C. 1999. The hunting apes: meat eating and the origins of human behavior. Princeton, NJ: Princeton University Press.
- Stanford CB, Bunn HT (eds.). 2001. *Meat eating and human evolution*. Oxford: Oxford University Press.
- Stearns SC. 1977. The evolution of life history traits: a critique of the theory and a review of the data. *Annu Rev Ecol Systemat* 8:145-171.
- Stearns SC. 1992. *The evolution of life histories*. Oxford: Oxford University Press.
- Stiner M. 2002. Carnivory, coevolution, and the geographic spread of genus *Homo*. *J Archaeol Res* 10:1-63.
- Sugiyama Y. 1994. Age-specific birth rate and life time reproductive success of chimpanzees at Bossou, Guinea. *Am J Primatol* 32:311-318.
- Sukumar R, Krishnamurthy V, Wemmer C, Rodden M. 1997. Demography of captive Asian elephants (*Elephas maximus*) in southern India *Zoo Biol* 16:263-272.
- Sutherland WJ, Grafen A, Harvey PH. 1986. Life history correlations and demography. *Nature* 320:88.
- Swisher C, Curtis G, Jacob T, Getty A, Suprijo A, Widiasmoro. 1994. Age of the earliest known hominids in Java. *Science* 263:118-121.
- Tardieu C. 1998. Short adolescence in early hominids: infantile and adolescent growth of the early human femur. *Am J Phys Anthropol* 107:163-178.
- Tice RR, Setlow RB. 1985. DNA repair and replication in aging organisms and cells. In: Finch CE, Schneider EL, editors. *Handbook of the biology of aging*, 2nd ed. New York: Van Nostrand. p 173-224.
- Tolmasoff JM, Ono T, Cutler RG. 1980. Superoxide dismutase: correlation with life span and specific metabolic rate in primate species. *Proc Natl Acad Sci USA* 77:2777-2781.
- Tooby J, DeVore I. 1987. The reconstruction of hominid behavioral evolution through strategic modeling. In: Kinzey W, editor. *The evolution of human behavior: primate models*. Albany, NY: SUNY Press. p 183-237.
- Treolar AE. 1981. Menstrual cyclicity and the premenopause *Maturitas* 3:249-264.
- Trinkaus E. 1986. The Neanderthals and modern human origins. *Annu Rev Anthropol* 15:198-218.
- Trinkaus E. 1995. Neanderthal mortality patterns. *J Archaeol Sci* 22:121-142.
- Trivers RL. 1972. Parental investment and sexual selection. In: Campbell B, editor. *Sexual selection and the descent of man*. Chicago: Aldine. p 139-179.
- Tuljapurkar S, Li N, Boe C. 2000. A universal pattern of mortality decline in the G7 countries. *Nature* 405:789-792.
- van Schaik CP. 1996. Social evolution in primates: the role of ecological factors and male behavior. *Proc Br Acad* 88:9-31.
- van Schaik CP, Janson C. 2000. *Infanticide by males and its implications*. Cambridge, UK: Cambridge University Press.
- van Schaik CP, Paul A. 1996. Male care in primates: does it ever reflect paternity? *Evol Anthropol* 5:152-156.
- Varki A. 2000. A chimpanzee genome project is a biomedical imperative. *Genome Res* 10:1065-1070.
- Vaupel JW, Carey JR, Christensen K, Johnson TE, Yashin AI, Holm NV, Iachine IA, Kannisto V, Khazaeli AA, Liedo P, Longo VD, Zeng Y, Manton KG, Curtsinger JW. 1998. Biodemographic trajectories of longevity. *Science* 280:855-860.
- Voland E, Biese J. 2002. Opposite effects of maternal and paternal grandmothers on infant survival in historical Krummhorn. *Behav Ecol Sociobiol* 52: 435-443.
- Vom Saal FS, Finch CE, Nelson JF. 1994. Natural history and mechanisms of reproductive aging in humans, laboratory rodents, and other selected vertebrates. In: Knobil E, Neill JD, editors. *The physiology of reproduction*, 2nd ed. New York: Raven Press. p 1213-1314.
- Walker LC. 1997. Animal models of cerebral $\beta$ -amyloid angiopathy. *Brain Res Rev* 25:70-74.
- Walker PL, Johnson JR, Lambert, PM. 1988. Age and sex bias in the preservation of human skeletal remains. *Am J Phys Anthropol* 76:183-188.
- Washburn SL. 1960. Tools and human evolution. *Sci Am* 203:63-75.
- Washburn SL, Avis V. 1958. Evolution and human behavior In: Roe A, Simpson GG, editors. *Behavior and evolution*. New Haven, CT: Yale University Press. p 421-436.
- Washburn SL, DeVore I. 1961. Social behavior of baboons and early man. In: Washburn S, editor. *Social life of early man*. Chicago: Aldine. p 91-105.
- Washburn SL, Lancaster CS. 1968. The evolution of hunting. In: Lee RB, DeVore I, editors. *Man the hunter*. Chicago: Aldine. p 293-303.
- Weiss KM. 1973. Demographic models for anthropology. *Soc Am Archaeol Mem* 27:1-86.
- Westendorp RGJ, Kirkwood TBL. 1998. Human longevity at the cost of reproductive success. *Nature* 396:743-746.
- Western D. 1979. Size, life history and ecology in mammals. *Afr J Ecol* 17:185-202.
- Williams GC. 1957. Pleiotropy, natural selection, and the evolution of senescence. *Evolution* 11:398-411.
- Williams GC. 1966a. Adaptation and natural selection: a critique of some current biological thought. Princeton, NJ: Princeton University Press.
- Williams GC. 1966b. Natural selection, the cost of reproduction, and a refinement of Lack's principle. *Am Nat* 199:687-690.
- Wolf ER. 1982. *Europe and the people without history*. Berkeley: University of California Press.
- Wood BA. 1992. Origin and evolution of the genus *Homo*. *Nature* 355:783-790.

- Wood JW. 1994. Dynamics of human reproduction. New York: Aldine de Gruyter.
- Wood BA, Collard M. 1999. The human genus. *Science* 284:65–71.
- Woodburn J. 1968. An introduction to Hadza ecology. In: Lee RB, DeVore I, editors. *Man the hunter*. Chicago: Aldine. p 49–55.
- Wrangham RW, Jones JH, Laden G, Pilbeam D, Conklin-Brittain NL. 1999. The raw and the stolen: cooking and the ecology of human origins. *Curr Anthropol* 40:567–594.
- Yu BP. 1996. Aging oxidative stress: modulation by dietary restriction. *Free Rad Biol Med* 21: 651–668.
- Zhu RK, Hoffman KA, Potts R, Deng CL, Pan YX, Guo B, Shi CD, Guo ZT, Yuan BY, Hou YM, Wang WW. 2001. Earliest presence of humans in northeast Asia. *Nature* 413:413–417.