

Grandmothering life histories and human pair bonding

James E. Coxworth^{a,b,c}, Peter S. Kim^d, John S. McQueen^d, and Kristen Hawkes^{a,1}

^aDepartment of Anthropology, University of Utah, Salt Lake City, UT 84112; ^bUtah Population Database, University of Utah, Salt Lake City, UT 84112; ^cAdvanced Solutions, Nielsen, Schaumburg, IL 60173; and ^dSchool of Mathematics and Statistics, University of Sydney, Sydney, NSW 2006, Australia

Contributed by Kristen Hawkes, August 4, 2015 (sent for review July 10, 2015; reviewed by Martin Daly)

The evolution of distinctively human life history and social organization is generally attributed to paternal provisioning based on pair bonds. Here we develop an alternative argument that connects the evolution of human pair bonds to the male-biased mating sex ratios that accompanied the evolution of human life history. We simulate an agent-based model of the grandmother hypothesis, compare simulated sex ratios to data on great apes and human hunter-gatherers, and note associations between a preponderance of males and mate guarding across taxa. Then we explore a recent model that highlights the importance of mating sex ratios for differences between birds and mammals and conclude that lessons for human evolution cannot ignore mammalian reproductive constraints. In contradiction to our claim that male-biased sex ratios are characteristically human, female-biased ratios are reported in some populations. We consider the likelihood that fertile men are undercounted and conclude that the mate-guarding hypothesis for human pair bonds gains strength from explicit links with our grandmothering life history.

grandmother hypothesis | human life history | human evolution | mate guarding | mating sex ratios

We call attention to evidence that connects the evolution of human pair bonds to the male-biased sex ratios in fertile ages that characterize human populations. As in mammals generally, age-specific mortality is higher in males than in females (e.g., refs. 1–3). However, this difference is overshadowed by a distinctive feature of human life history: Oldest ages at parturition are about the same in humans as in other living hominids, the great apes (4, 5), whereas longevity is substantially greater and male fertility continues to older ages (6). Exceptional longevity with a distinctive postmenopausal life stage (7–9) may have evolved in our lineage when grandmothers' subsidies for weaned dependents allowed mothers to have next babies sooner. According to this grandmother hypothesis (10–16), longevity increased as longer-lived grandmothers could help more and so left more longer-lived descendants of both sexes. Women's postfertile life stage (7) produces a bias in the sex ratio of fertile adults with repercussions for male strategies. As longevity increased, older-aged males expanded the pool of competitors for the still-fertile females. With more competitors for each paternity, males' average success in finding new mates inevitably declined until defending a current mate became the better option. Our distinctive life history thus supplies previously unrecognized support for a mate-guarding hypothesis for the evolution of human pair bonds.

Here we simulate hominid mating sex ratios with an agent-based model of the evolution of human longevity via grandmothering (13, 15). We then compare simulated sex ratios to demographic data from both great apes and human hunter-gatherers. Having identified the human bias, we connect it to increased male payoffs for mate guarding, noting some broad patterns in humans, the tradeoffs observed in other taxa, and a history of modeling in which increased guarding is the likely outcome of more competing males.

We then consider a recent model (17) of the evolution of sex roles built to show that anisogamy, the primary sex difference of large (female) and small (male) gametes (18), is insufficient by itself to explain why females care for offspring more often whereas males more often compete for mates. Mating sex ratios

are decisive for mating strategies in this model. Although it does not include guarding as an option, it does highlight a connection between mating sex ratios and broad differences between mammals and birds. We elaborate key phylogenetic constraints, consider recent work on birds, and underline features of mammalian offspring production that temper direct application of the model to mammals, including hominids.

An important challenge to our claim that human life history entails male-biased mating sex ratios comes from reports of female bias in some human populations. We identify common measurement problems and link men's age-specific fertilities to status hierarchies, concluding that the mate-guarding hypothesis remains both promising and directly relevant to explaining the long-recognized importance of male alliances in human societies.

Grandmothering Simulations

We track mating sex ratios through simulations of an agent-based model first built to investigate the evolution of human longevity via grandmothering (13, 15). For reasons elaborated below, we follow both the adult sex ratio (ASR), defined as the ratio of males to females in the fertile ages, and the operational sex ratio (OSR), which counts only the subset of adults currently capable of a conception (19) (see *Supporting Information* for model parameters and definitions).

Elsewhere we have shown that simulations of this model result in two life history equilibria: a great ape-like one with no grandmothering and a human hunter-gatherer-like one when grandmothers' subsidies allow mothers to have next babies sooner (15). Not surprisingly, each equilibrium is associated with distinct, characteristic sex ratios in the fertile ages (Fig. 1). Of 30 simulations without grandmothering run over a million years, the ASR (Fig. 1A) fluctuates around an average of 0.77 males for every female whereas OSR (Fig. 1C) remains at about 50.

Significance

Pair bonds are universal in human societies and distinguish us from our closest living relatives. They characteristically involve men's proprietary claims over women—mate guarding—which in animals generally is both predicted and observed to be more frequent when sex ratios in the fertile ages are male-biased. A marked male bias in the fertile ages evolved in our lineage as longevity increased without an extension of female fertility. We compare the sex-ratio shift in simulations of the grandmother hypothesis to demographic data from chimpanzees and human hunter-gatherers then connect the expanded proportions of older men to benefits for mate guarding, the evolution of pair bonds, and the long recognized importance of male alliances in human social life.

Author contributions: J.E.C. and K.H. designed research; J.E.C. and P.S.K. performed research; J.S.M. contributed new reagents/analytic tools; J.E.C. and P.S.K. analyzed data; and J.E.C., P.S.K., and K.H. wrote the paper.

Reviewers included: M.D., McMaster University.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. Email: hawkes@anthro.utah.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1599993112/-DCSupplemental.




Fig. 1. Time evolutions of ASRs and OSRs with and without grandmothering. (A) ASRs of 30 simulations over 1 million y without grandmothering. Each simulation is shown in light gray. The average of the 30 simulations is shown in black and ends at an ASR of 0.77. The ending point of the simulation shown in medium gray serves as the starting point for the 30 new simulations with grandmothering shown in B. (B) ASRs of 30 simulations over 2 million y with grandmothering. Each simulation is shown in gray. The average of the 30 simulations, in black, ends at an ASR of 1.56. (C) OSRs of 30 simulations over 1 million y without grandmothering. Colors as in A. The average of the 30 simulations ends at an OSR of 50. (D) OSRs of 30 simulations over 2 million y with grandmothering. Colors as in B. The average ends at an OSR of 111.

To evaluate the effect of grandmothering on the mating sex ratios, we use the end point of a simulation chosen at random to start 30 new simulations with grandmothering. All 30 simulations evolve away from the previously stable ape-like mating sex ratios toward values roughly two times higher. ASR (Fig. 1B), initially female-biased, increases with grandmothering to about 1.56 fertile males for every female in the fertile ages, whereas OSR (Fig. 1D) rises to ~111 available males for every female ready to conceive.

Life Tables

To see whether the simulated values are empirically plausible we compare them to the same sex ratios calculated using life tables for living populations. Our empirical comparison for the simulated ape-like equilibrium is limited to chimpanzees because they are the only apes for which life tables based on multiple, long-studied populations have been published. Bonobo population dynamics are still poorly known. We expect, based on arguments here, ASRs to be higher in mate-guarding gorillas than in chimpanzees. Demography of the Virunga population (20) indicates that, unlike chimpanzees, mortality is not higher in males until the oldest ages. Sexes are not always distinguished in more recent gorilla censuses (21, 22), so we await more data. For orangutans (23), relatively low male mortality is also indicated, but it is difficult to know which males to include among the competitors for mates.

We estimate chimpanzee mating sex ratios using demographic data from the five-study site synthesis of Hill et al. (24) and also include Muller and Wrangham's (25) more recent (though very small) sample for one of those sites where overall mortality is lowest.

For humans, age structure data come from four well-known hunter-gatherer groups: Howell's (26, 27) life tables for Dobe !Kung, Hill and Hurtado's (28) for forest Ache, Hill et al.'s (29) for Hiwi, and Blurton Jones's (30) for Hadza. We chose these groups because of their reliance on wild foods and the quality of demographic data collected among them. Although contemporary foragers are not living fossils, their mortality regimes make demographic patterns observed among them the best representation of human demography during our evolutionary past.

Table 1 shows that the empirical values computed from chimpanzee sources (ASRs of 0.47 and 0.70; OSRs of 41 and 61) are in general accord with the simulated values without grandmothering. Among the hunter-gatherers (Table 2) ASRs average 1.64, whereas OSRs average 83 (range 69–92), mirroring the equilibrium ASR and OSR values reached in our simulations with grandmothering. Two attributes of these results are especially important. First, every empirical human ASR and OSR is encompassed, or nearly so, in every simulation. Second, in both the real world and simulations with grandmothering the human ratios are distinctly higher than those of the great apes.

Mating Sex Ratios and Mating Strategies

Marlowe and Berbesque (37) pointed to the effects of a strongly male-biased OSR on male competition and marriage resulting from women's long postfertile life stage. In general, male responses to such a situation should depend on the costs and benefits of available alternatives. Parker (38) was the first to suggest a positive relationship between the relative number of males in a population and the fitness benefits of defending females, or mate guarding. Predictions about the evolution of mate guarding have been elaborated by a number of subsequent

Table 1. Demographic parameters for chimpanzees

Population	Males age 15+ years	Females age 10–45 years	Birth interval, year	Male paternity, days/year	Females fecundable days per cycle	Cycles to conception	ASR M/F	OSR M/F
Synthesis of five (24)	0.321	0.679	5.72	365	6	4	0.47	40.13
Kanyawara (25)	0.411	0.589	5.72	365	6	4	0.70	60.70

Assuming stationary populations, the mortality curve mirrors the age structure. To model age structures we used probability of survival to each age in the published life tables, summing the calculated number of survivors for males and females to each of the fertile ages, then dividing the sum for each sex by their combined total to get the fraction fertile adults by sex (columns 2 and 3). We included males older than 15 years and females between ages 10 and 45 years (31). Data on chimpanzee birth intervals come from averaging reports in Knott (32).

For OSR we used the formula for nonseasonal breeders derived by Mitani et al. (33), where

$$OSR = \frac{m \cdot B \cdot 365}{f \cdot \sum_{i=1}^n s_i}$$

In this equation, m and f are the fraction of fertile adults that are male and female, respectively; B is the average birth interval; and 365 is the days per year that males can compete for a paternity. The summation in the denominator is the fecundable days per birth interval for fertile females. It depends on the number of conception risk days per estrous cycle (s) and the number of cycles per conception (n). As in the simulations we use observations from humans (34, 35) and fix $s = 6$ and $n = 4$.

treatments (e.g., refs. 39–44). The hypothesized relationship between biased sex ratios and mate guarding has also been supported by observational and experimental studies conducted across a range of species (45–50).

Although mate guarding is not synonymous with pair bonding, similar tradeoffs are involved. Guarding a mate (or potential mate) is beneficial for members of the more numerous sex when it leads to greater reproductive success (RS) than continually seeking out new mates (38). The logic underlying pair bonding is the same: individuals of the more prevalent sex should guard continuously (i.e., form pairs) when doing so leads to greater RS than the alternatives (44).

Parker and Stuart's (39) model assumed that males acquire females through either guarding a current mate (G) or searching for another one (S). Clutton-Brock and Parker (51) retained this S vs. G notation, but they posed the alternatives differently (ref. 51, p. 447), defining "time in" the mating pool (S) to include both searching and guarding, with guarding also part of "time out" (G).

Of special relevance here, Kokko and Jennions (17) used Clutton-Brock and Parker's time in vs. time out framework to build new models for the evolution of mating behavior. Revisiting critiques of Trivers' (1) influential arguments about the role of parental investment (52–54), they constructed a model to show "from first principles" why anisogamy alone cannot explain sex-role divergence. Then they faced the "more fundamental question: what factors create the asymmetry that biases females towards caring for offspring and males towards competing for mates?" (ref. 17, p. 920). Their answer followed Queller's (53) earlier analysis that identified multiple mating by females, which

lowers average male parentage and so favors less parental care from males, and sexual selection, which makes expected mating success higher than average for some males.

The particular importance of ASR (54–57) was further emphasized by Kokko and Jennions (17). Their model did not include a mate-guarding option but it has influenced subsequent discussion of the effects of human mating sex ratios on male strategies. In their model individuals can enter the mating pool or stay out to gain benefits in offspring survival from continued parental care. Because the sex that is more numerous must have a lower average mating rate, that sex does better to stay out longer. As they do, bias in OSR declines toward parity and sex roles diverge less. How even the OSR can get depends on the ASR because of the Fisher condition (55–58); because everyone has one mother and one father, the number of offspring produced by one sex must be the same as the number produced by the other. This model has been read by some (59) as undermining OSR as a predictor of mating patterns, warranting focus on ASR instead.

If the tradeoffs for hominid males were the ones assumed in Kokko and Jennions' (17) basic model, the evolution of increasingly male-biased OSRs would be unstable. Where OSR is male-biased, males take longer to find another mate and so do better to choose an alternative to searching for them. While emphasizing this important effect, the only alternative considered in their basic model is to supply parental care, with each additional unit of care (from either sex) earning equivalent marginal gains in offspring survival. Given those assumptions, male bias in the mating pool must decline as males stay out to parent longer.

Table 2. Demographic parameters for human hunter-gatherers

Population	Males age 20–65 years	Females age 20–40 years	Birth interval, year	Male paternity, days/year	Female fecundable days per cycle	Cycles to conception	ASR M/F	OSR M/F
Dobe !Kung (26, 27)	0.593	0.407	4.17	365	6	4	1.46	92.40
Ache forest (28)	0.652	0.348	2.44	365	6	4	1.87	69.52
Hiwi (29)	0.618	0.382	3.70	365	6	4	1.62	91.04
Hadza (30)	0.616	0.384	3.23	365	6	4	1.60	78.80

Assuming stationary populations, the mortality curve mirrors the age structure. To model age structures we used probability of survival to each age in the published life tables, summing the calculated number of survivors for men and women to each of the fertile ages, then dividing the sum for each sex by their combined total to get the fraction fertile adults by sex (columns 2 and 3). We included men from 20 and 65 years based on reported age ranges of fertilities from the ethnographers and those reported by Tuljapurkar et al. (36). Women from 20 to 40 years are included based on average ages of first and last birth (4). Since populations are younger when increasing, and the growth rates for the human populations range from 0.26% to 2.5%, our figures overestimate the ASRs by 2–12%.

We computed birth intervals by dividing years of female fertility by the total fertility rate reported in the ethnographic sources. OSR was calculated as in Table 1.

However, strongly male-biased OSRs persist across the entire class of mammals where male parental care is rare. In our simulations and reference populations the OSRs are male-biased throughout, as is typical of mammals. Even though birth intervals are shorter in humans, the addition of older males makes the bias twice as strong in humans as in chimpanzees. However, the non-human ASR is female-biased (the common mammal pattern), while—again because of so many old men—the human ASR is male-biased—a pattern not typical of mammals but of birds. Kokko and Jennions (17) refer to this broad taxonomic difference in ASRs and relate it to the sex roles generally recognized as typical of these two classes of vertebrates. Do the “bird-like” human ASRs indicate useful parallels with class Aves (19, 60–62)?

Key Constraints

In birds either males or females can brood and guard eggs, as well as guard and feed nestlings. OSR depends on how much each sex is occupied with care. High frequency of biparental care (60) implies that sex-role convergence is common. Especially striking evidence of this near sexual equivalence comes from penduline tits (*Remiz pendulinus*), where either the male or the female always deserts to mate again and nearly a third of nests fail from desertion by both parents (63). Liker et al. (64) found that in shore birds (where sex-role reversal is common), the variation in sex roles correlates with ASRs as predicted (17, 56) and proposed (ref. 64, p. 3) “that the evolutionary flexibility of both sexes to provide full care on their own and variation in ASR among species are among the key factors that facilitate the evolution of diverse sex roles.” Extending the analysis, they found ASRs predict how much males care among a wider range of wild bird populations (65).

Even in birds, however, the sexes cannot make equivalent contributions to offspring production (66). The difference is much larger in mammals. That difference, as Kokko and Jennions (ref. 17, p. 940) note, “might be the result of simple constraint (e.g. that the evolution of lactation is difficult for male mammals).” Once internal gestation and lactation have evolved, care is much less interchangeable between the sexes. Fundamental mammalian constraints apply to the primate radiation and are directly relevant to expectations we develop about male-biased OSRs and ASRs in our lineage.

Schacht and colleagues (ref. 59, p. 215) downplay this issue to argue that emphasis on OSR as a predictor of mating strategies has been misleading because, “contrary to the intuitions drawn from Emlen and Oring 1977, a male-biased OSR only accurately predicts intense sexual selection among males under a limited set of circumstances. . . .” For this they cite Kokko et al. (67), who note that although Emlen and Oring’s (19) emphasis on OSR as an index of mating behavior seems borne out empirically, the ties between OSR and sexual selection depend on “conditions that make selection favour traits that reduce the time it takes to acquire another mating” (ref. 67, p. 1341). When time out is a necessarily large fraction of a fertile lifespan, higher mating rates make little difference to lifetime RS (ref. 67, p. 1349). This is precisely the case for female mammals and especially for female primates, who are committed by gestation and long lactation to extended times out.

In contrast, mating rates do make a difference to lifetime RS for mammalian males. When too many males lower the expected success rate for seeking a new mate, doing something else may be a better option. Harts and Kokko (44) investigated mate guarding as the alternative and showed “in line with earlier work more male competitors select for more guarding” (ref. 44, p. 2842). Among primates with multilevel social systems (68–70), including humans (71), a male’s RS usually depends on his success at claiming and retaining mates, which depends on other males deferring to his claims.

Measuring Human Sex Ratios: ASRs or Elder Advantage?

In our simulations mating sex ratios become increasingly male-biased with a growing fraction of elders as a human-like life history evolves from a great ape-like one. This reflects the approximate doubling of longevity while female fertility maintains its ancestral decline to near zero by 45. Although mortality is generally higher in men than in women (2, 3), the increased number of older men who are not yet frail makes human ASRs inevitably male-biased.

How, then, can some explorations of human mating sex ratios report so many to be female-biased? Using variance in RS reported by Brown et al. (72), Kokko and Jennions (ref. 73, pp. 113–114) used the Fisher condition and the RSs reported to infer ASRs for 18 human populations, estimating them to be about even for the societies classified as monogamous or serially monogamous and significantly female-biased for the groups classified as polygynous. For nine cases Brown et al. (72) classified as polygynous, the mean RS for males averages 31% higher than the female mean—possibly due, as Kokko and Jennions (73) conclude, to female-biased ASRs. But Brown et al. (ref. 72, p. 300) note another possibility: missing men who were less successful in the mating competition.

An earlier study by Ember (74) also found an association between polygyny and female-biased sex ratios using data from the Human Relations Area Files. While high enough male mortality could make ASRs female-biased, Ember’s sex ratios were not ASRs but sex ratios of whole populations—immatures and postfertile women included. Schacht et al. (ref. 59, p. 218) note this difficulty with “which sex ratio.” They review social science research into sex-ratio effects listing 20 studies in their first table; none of them used fertile ages.

Figure 2 in Schacht et al. (ref. 59, p. 217) is the most serious challenge to our characterization of human mating sex ratios as male-biased. It plots results from the authors’ queries to ethnographers about 15 traditional populations. Here, they say, “Sex ratio is determined from the ethnographers’ data on the number of individuals of mating age in their population, hence it approximates ASR.” Eight of these societies—more than half—have female-biased ASRs, a pattern that our model and arguments suggest is “not human.”

At least three factors could lead ethnographers to underestimate sex ratios in the fertile ages. First, physiologically fertile young men may be excluded because local conventions class them as not yet of mating age. Székely et al. (ref. 75, p. 1501), reviewing broader cross-species issues of ASR and OSR associations with breeding systems, define an adult as “an animal (male or female) that is physiologically capable of producing offspring.”

Second, counting unmarried men is difficult if they move more often than others, a common pattern among hunter-gatherers. Such high mobility gave Hawkes et al. (ref. 76, p. 683) an insufficient sample of unmarried Hadza men to compare their time allocation patterns with those of married men. Székely et al. (ref. 75, p. 1501) note similar ascertainment biases with other animals, because “nonbreeding adults (e.g. floaters) often . . . remain unnoticed.”

A third, closely related possibility is that ethnographers chose not to include unmarried men. For example, polygynous Kipsigis have a notably low ASR. Reporting that Kipsigis men have much higher average RS compared with women (which, given the Fisher condition, would require extremely high male mortality), Borgerhoff Mulder (ref. 77, p. 433) said that her tabulation excluded “poor men, who leave the community to become plantation laborers and marry late in life, if at all.”

Marlowe and Berbesque (37) showed the advantage that elder men can have in ethnographically known hunter-gatherers by plotting the relationship between age of first marriage and the extent of polygyny across 130 hunter-gatherer societies. Their figure 1 (ref. 37, p. 836) shows that men’s age at first marriage

36. Tuljapour SD, Puleston CO, Gurven MD (2007) Why men matter: Mating patterns drive evolution of human lifespan. *PLoS One* 2(8):e785.
37. Marlowe FW, Berbesque JC (2012) The human operational sex ratio: Effects of marriage, concealed ovulation, and menopause on mate competition. *J Hum Evol* 63(6):834–842.
38. Parker GA (1974) Courtship persistence and female-guarding as male time investment strategies. *Behaviour* 48(1–2):157–184.
39. Parker GA, Stuart RA (1976) Animal behavior as a strategy optimizer: Evolution of resource assessment strategies and optimal emigration thresholds. *Am Nat* 110(976):1055–1076.
40. Grafen A, Ridley M (1983) A model of mate guarding. *J Theor Biol* 102(4):549–567.
41. Hawkes K, Rogers AR, Charnov EL (1995) The male's dilemma: Increased offspring production is more paternity to steal. *Evol Ecol* 9:662–677.
42. Yamamura N, Tsuji N (1989) Postcopulatory guarding strategy in a finite mating period. *Theor Popul Biol* 35:36–50.
43. Fromhage L, Elgar MA, Schneider JM (2005) Faithful without care: The evolution of monogyny. *Evolution* 59(7):1400–1405.
44. Harts AMF, Kokko H (2013) Understanding promiscuity: When is seeking additional mates better than guarding an already found one? *Evolution* 67(10):2838–2848.
45. Møller AP (1987) Extent and duration of mate guarding in swallows *Hirundo rustica*. *Ornis Scand* 18(2):95–100.
46. Rondeau A, Sainte-Marie B (2001) Variable mate-guarding time and sperm allocation by male snow crabs (*Chionoecetes opilio*) in response to sexual competition, and their impact on the mating success of females. *Biol Bull* 201(2):204–217.
47. Sherman PW (1989) Mate guarding as paternity insurance in Idaho ground squirrels. *Nature* 338(6214):418–420.
48. Alcock J (1994) Postinsemination associations between males and females in insects: the mate-guarding hypothesis. *Annu Rev Entomol* 39:1–21.
49. Jormalainen V (1998) Precopulatory mate guarding in crustaceans: Male competitive strategy and intersexual conflict. *Q Rev Biol* 73(3):275–304.
50. Weir LK, Grant JWA, Hutchings JA (2011) The influence of operational sex ratio on the intensity of competition for mates. *Am Nat* 177(2):167–176.
51. Clutton-Brock TH, Parker GA (1992) Potential reproductive rates and the operation of sexual selection. *Q Rev Biol* 67(4):437–456.
52. Dawkins R, Carlisle TR (1976) Parental investment, mate desertion and a fallacy. *Nature* 262:131–133.
53. Queller DC (1997) Why do females care more than males? *Proc Biol Sci* 264:1555–1557.
54. Wade MJ, Shuster SM (2002) The evolution of parental care in the context of sexual selection: A critical reassessment of parental investment theory. *Am Nat* 160(3):285–292.
55. Houston AI, McNamara JM (2002) A self-consistent approach to paternity and parental effort. *Philos Trans R Soc Lond B Biol Sci* 357(1419):351–362.
56. McNamara JM, Székely T, Webb JN, Houston AI (2000) A dynamic game-theoretic model of parental care. *J Theor Biol* 205(4):605–623.
57. Székely T, Webb JN, Cuthill IC (2000) Mating patterns, sexual selection and parental care: An integrative approach. *Vertebrate Mating Systems*, eds Apollonio M, Festa-Bianchet M, Mainardi D (World Science, London), pp 194–223.
58. Fisher RA (1930) *The Genetical Theory of Natural Selection* (Oxford Univ Press, Oxford).
59. Schacht R, Rauch KL, Borgerhoff Mulder M (2014) Too many men: The violence problem? *Trends Ecol Evol* 29(4):214–222.
60. Lack D (1968) *Ecological Adaptations for Breeding in Birds* (Methuen, London).
61. Birkhead T (1987) Sperm competition in birds. *Trends Ecol Evol* 2(9):268–272.
62. Westneat DF, Stewart IRK (2003) Extra-pair paternity in birds: Causes, correlates, and conflict. *Annu Rev Ecol Syst* 34:365–396.
63. Szentirmai I, Székely T, Komdeur J (2007) Sexual conflict over care: Antagonistic effects of clutch desertion on reproductive success of male and female penduline tits. *J Evol Biol* 20(5):1739–1744.
64. Liker A, Freckleton RP, Székely T (2013) The evolution of sex roles in birds is related to adult sex ratio. *Nat Commun* 4:1587.
65. Liker A, Freckleton RP, Székely T (2014) Divorce and infidelity are associated with skewed adult sex ratios in birds. *Curr Biol* 24(8):880–884.
66. Sibly RM, et al. (2012) Energetics, lifestyle, and reproduction in birds. *Proc Natl Acad Sci USA* 109(27):10937–10941.
67. Kokko H, Klug H, Jennions MD (2012) Unifying cornerstones of sexual selection: operational sex ratio, Bateman gradient and the scope for competitive investment. *Ecol Lett* 15(11):1340–1351.
68. Kummer H (1968) *Social Organization of Hamadryas Baboons: A Field Study* (Univ of Chicago Press, Chicago).
69. Swedell L, Leedom L, Saunders J, Pines M (2013) Sexual conflict in a polygynous primate: Costs and benefits of a male-imposed mating system. *Behav Ecol Sociobiol* 68:263–273.
70. Grueter CC, Chapais B, Zinner D (2012) Evolution of multilevel social systems in nonhuman primates and humans. *Int J Primatol* 33(5):1002–1037.
71. Rodseth L (2013) From bachelor threat to fraternal security: Male associations and modular organization in human societies. *Int J Primatol* 33:1194–1214.
72. Brown GR, Laland KN, Borgerhoff Mulder M (2009) Bateman's principles and human sex roles. *Trends Ecol Evol* 24(6):297–304.
73. Kokko H, Jennions MD (2012) Sex differences in parental care. *The Evolution of Parental Care*, eds Royle NJ, Smiseth PT, Kolliker M (Oxford Univ Press, Oxford), pp 110–116.
74. Ember M (1974) Warfare, sex ratio, and polygyny. *Ethnology* 13(2):197–206.
75. Székely T, Weissing FJ, Komdeur J (2014) Adult sex ratio variation: Implications for breeding system evolution. *J Evol Biol* 27(8):1500–1512.
76. Hawkes K, O'Connell JF, Blurton Jones NG (1997) Hadza women's time allocation, offspring provisioning and the evolution of post-menopausal lifespans. *Curr Anthropol* 38(4):551–577.
77. Borgerhoff Mulder M (1988) Reproductive success in three Kipsigis cohorts. *Reproductive Success: Studies of Individual Variation in Contrasting Breeding Systems*, ed Clutton-Brock TH (Univ of Chicago Press, Chicago), pp 419–435.
78. Darwin C (1871) *The Descent of Man, and Selection in Relation to Sex* (Murray, London); reprinted (1981) (Princeton Univ Press, Princeton).
79. Collier JF, Rosaldo MZ (1981) Politics and gender in simple societies. *Sexual Meanings: The Cultural Construction of Gender and Sexuality*, eds Ortner SB, Whitehead H (Cambridge Univ Press, Cambridge, UK), pp 275–329.
80. Vinicius L, Mace R, Miglano A (2014) Variation in male reproductive longevity across traditional societies. *PLoS One* 9(11):e112236.
81. Washburn SL, Lancaster CS (1968) The evolution of hunting. *Man the Hunter*, eds Lee RB, DeVore I (Aldine, New York), pp 293–303.
82. Isaac G (1978) The food-sharing behavior of protohuman hominids. *Sci Am* 238(4):90–108.
83. Lovejoy CO (1981) The origin of man. *Science* 211(4480):341–350.
84. Fisher H (1992) *The Anatomy of Love* (Norton, New York).
85. Kaplan H, Hill K, Lancaster J, Hurtado AM (2000) A theory of human life history evolution: Diet, intelligence, and longevity. *Evol Anthropol* 9:156–185.
86. Kaplan H, Gurven M, Winking J, Hooper PL, Stieglitz J (2010) Learning, menopause, and the human adaptive complex. *Ann N Y Acad Sci* 1204:30–42.
87. Wilson M, Daly M (1992) The man who mistook his wife for a chattel. *The Adapted Mind: Evolutionary Psychology and the Generation of Culture*, eds Barkow JH, Cosmides L, Tooby J (Oxford Univ Press, New York), pp 289–322.
88. Hawkes K (2004) Mating, parenting and the evolution of human pair bonds. *Kinship and Behavior in Primates*, eds Chapais B, Berman C (Oxford Univ Press, Oxford), pp 443–473.
89. Chapais B (2008) *Primeval Kinship: How Pairbonding Gave Birth to Human Society* (Harvard Univ Press, Cambridge, MA).
90. Chapais B (2010) The deep structure of human society: Primate origins and evolution. *Mind the Gap: Tracing the Origins of Human Universals*, eds Kappeler PM, Silk JB (Springer, Berlin), pp 19–51.
91. Komers PE, Brotherton PN (1997) Female space use is the best predictor of monogamy in mammals. *Proc Biol Sci* 264(1386):1261–1270.
92. van Shaik CP, Kappeler PM (2003) The evolution of social monogamy in mammals. *Monogamy: Mating Strategies and Partnerships in Birds, Humans, and Other Mammals*, eds Reichard UH, Boesch C (Cambridge Univ Press, Cambridge, UK), pp 59–80.
93. Lukas D, Clutton-Brock TH (2013) The evolution of social monogamy in mammals. *Science* 341(6145):526–530.
94. Blurton Jones NG, Marlowe FW, Hawkes K, O'Connell JF (2000) Paternal investment and hunter-gatherer divorce rates. *Adaptation and Human Behavior: An Anthropological Perspective*, eds Cronk L, Chagnon N, Irons W (Aldine de Gruyter, New York), pp 31–55.
95. Guttentag M, Secord PF (1983) *Too Many Women? The Sex Ratio Question* (Sage, Beverly Hills, CA).
96. Langness LL (1967) Sexual antagonism in the New Guinea Highlands: A Bena Bena example. *Oceania* 37(3):161–177.
97. Whiting JWM, Whiting BB (1975) Aloofness and intimacy of husbands and wives: A cross-cultural study. *Ethos* 3(2):183–207.
98. Smuts BB (1992) Male aggression against women: An evolutionary perspective. *Hum Nat* 3(1):1–44.
99. Charnov EL (1991) Evolution of life history variation among female mammals. *Proc Natl Acad Sci USA* 88(4):1134–1137.
100. Charnov EL (1993) *Life History Invariants: Some Explorations of Symmetry in Evolutionary Ecology* (Oxford Univ Press, Oxford).
101. Boesch C, Bolé C, Eckhardt N, Boesch H (2010) Altruism in forest chimpanzees: The case of adoption. *PLoS One* 5(1):e8901.
102. Sear R, Mace R (2008) Who keeps children alive? A review of the effects of kin on child survival. *Evol Hum Behav* 29(1):1–18.